

36-2048: Anti-CFTR (Cystic Fibrosis Transmembrane Conductance Regulator) Monoclonal Antibody (Clone: CFTR/1785)

| | |
|--------------------------------|--|
| Clonality : | Monoclonal |
| Clone Name : | CFTR/1785 |
| Application : | IHC |
| Reactivity : | Human |
| Gene : | CFTR |
| Gene ID : | 1080 |
| Uniprot ID : | P13569 |
| Alternative Name : | ABC35; ATP Binding Cassette Superfamily C Member 7 (ABCC7); cAMP-dependent chloride channel; CFTR; CFTR/MRP; Channel conductance-controlling ATPase; Cystic Fibrosis Transmembrane Conductance Regulator; MRP7; TNR CFTR |
| Isotype : | Mouse IgG2b, kappa |
| Immunogen Information : | Recombinant fragment (around aa 258-385) of human CFTR protein (exact sequence is proprietary) |

Description

Recognizes a protein of 165-170kDa, identified as cystic fibrosis transmembrane conductance regulator (CFTR). CFTR is composed of two membrane-spanning domains (MSD), two nucleotide-binding domains (NBD), and an R domain. It is structurally similar to multidrug resistance (Mdr1) protein and both are members of the superfamily of ATP-binding cassette (ABC) transporters, also known as traffic ATPases, which are implicated in the movement of various substrates. The CFTR protein is a small conductance adenosine 3',5'-cyclic monophosphate (cAMP)-activated chloride ion channel found in the apical membranes of epithelia within the pancreas, airway, intestine, bile duct, sweat gland, and male genital ducts. CFTR is a valuable marker of human pancreatic duct cell development and differentiation.

Product Info

| | |
|----------------------------|--|
| Amount : | 20 µg / 100 µg |
| Content : | 200µg/ml of Ab Purified from Bioreactor Concentrate by Protein A/G. Prepared in 10mM PBS with 0.05% BSA & 0.05% azide. Also available WITHOUT BSA & azide at 1.0mg/ml. |
| Storage condition : | Antibody with azide - store at 2 to 8°C. Antibody without azide - store at -20 to -80°C. Antibody is stable for 24 months. |

Application Note

Immunohistochemistry (Formalin-fixed) (1-2ug/ml for 30 minutes at RT)(Staining of formalin-fixed tissues is enhanced by heating tissue sections in 10mM Tris with 1mM EDTA, pH 9.0 for 45 min at 95°C followed by cooling at RT for 20 minutes)




Fig.1: Formalin-fixed, paraffin-embedded human Pancreas stained with CFTR Mouse Monoclonal Antibody (CFTR/1785).




Fig. 2: SDS-PAGE Analysis Purified CFTR Mouse Monoclonal Antibody (CFTR/1785).