

## 36-2075: Anti-Chromogranin A / CHGA (Neuroendocrine Marker) Monoclonal Antibody (Clone: CHGA/777)-CF488

|                                |   |
|--------------------------------|---|
| <b>Clonality :</b>             | Monoclonal  |
| <b>Clone Name :</b>            | CHGA/777  |
| <b>Application :</b>           | FACS,IF   |
| <b>Reactivity :</b>            | Human   |
| <b>Conjugate :</b>             | CF488   |
| <b>Gene :</b>                  | CHGA  |
| <b>Gene ID :</b>               | 1113  |
| <b>Uniprot ID :</b>            | P10645  |
| <b>Alternative Name :</b>      | Beta-Granin; CGA; CHGA; Chromogranin A Parathyroid Secretory Protein 1; ER-37; Pancreastatin; Parastatin; Pituitary Secretory Protein I; SP-I; Vasostatin I or II |
| <b>Isotype :</b>               | Mouse IgG1, kappa   |
| <b>Immunogen Information :</b> | Recombinant human CHGA protein  |

### Description

Chromogranin A is present in neuroendocrine cells throughout the body, including the neuroendocrine cells of the large and small intestine, adrenal medulla and pancreatic islets. It is an excellent marker for carcinoid tumors, pheochromocytomas, paragangliomas, and other neuroendocrine tumors. Co-expression of chromogranin A and neuron specific enolase (NSE) is common in neuroendocrine neoplasms. Reportedly, co-expression of certain keratins and chromogranin indicates neuroendocrine lineage. The presence of strong anti-chromogranin staining and absence of anti-keratin staining should raise the possibility of paraganglioma. The co-expression of chromogranin and NSE is typical of neuroendocrine neoplasms. Most pituitary adenomas and prolactinomas readily express chromogranin A.

### Product Info

|                            |  |
|----------------------------|--|
| <b>Amount :</b>            | 0.5 ml at 100µg/ml   |
| <b>Content :</b>           | Antibody Purified from Bioreactor Concentrate by Protein A/G and conjugated to various reporter molecules. Prepared in 10mM PBS with 0.05% BSA and 0.05% azide. Contact us if you require this Ab in a different format. |
| <b>Storage condition :</b> | Antibody with azide - store at 2 to 8°C. Antibody without azide - store at -20 to -80°C. Antibody is stable for 24 months.   |

### Application Note

Flow Cytometry (5ul per test per one million cells or 5ul per 100ul of whole blood);Immunofluorescence (1:50-1:100);