

36-2116: Anti-CD21 (Mature B-Cell & Follicular Dendritic Cell Marker) Monoclonal Antibody(Clone: CR2/2754)

| | |
|--------------------------------|--|
| Clonality : | Monoclonal |
| Clone Name : | CR2/2754 |
| Application : | ELISA,FACS,WB,IF,IHC |
| Reactivity : | Human |
| Gene : | CR2 |
| Gene ID : | 1380 |
| Uniprot ID : | P20023 |
| Alternative Name : | CD21; Complement C3d receptor (C3DR); Complement Receptor type 2 (CR2); CVID7; EBV receptor; EBV-R; Epstein-Barr virus receptor; EVBR; SLEB9 |
| Isotype : | Mouse IgG1, kappa |
| Immunogen Information : | Recombinant fragment (around aa 142-240) of human CR2 (CD21) protein (exact sequence is proprietary) |

Description

Recognizes a protein of 140kDa, which is identified as the complement receptor 2 (CR2) or CD21. This protein is expressed strongly on mature B cells, follicular dendritic cells and weakly on immature thymocytes and T lymphocytes. In B-cell ontogeny, CD21 appears after the pre-B-stage, is maintained during peripheral B-cell development and is lost upon terminal differentiation into plasma cells. CD21 expression is also gradually lost after stimulation of B cells in vitro. CD21 functions as receptor for C3d, C3dg and iC3b Complement components, for EBV and for IFNalpha. CD21 binds to CD23 and associates with CD19, CD81 and Leu13 to form a large signal-transduction complex involved in B cell activation.

Product Info

| | |
|----------------------------|---|
| Amount : | 20 µg / 100 µg |
| Content : | 200 µg/ml of Ab Purified from Bioreactor Concentrate by Protein A/G. Prepared in 10mM PBS with 0.05% BSA & 0.05% azide. Also available WITHOUT BSA & azide at 1.0mg/ml. |
| Storage condition : | Antibody with azide - store at 2 to 8°C. Antibody without azide - store at -20 to -80°C. Antibody is stable for 24 months. Non-hazardous. |

Application Note

ELISA (For coating, order antibody without BSA);,Flow Cytometry (1-2ug/million cells); Cells are fixed using 3% PFA.,Western Blot (1-2ug/ml);,Immunofluorescence (1-2ug/ml);,Immunohistology (Formalin-fixed) (1-2ug/ml for 30 min at RT),(Staining of formalin-fixed tissues requires boiling tissue sections in 10mM Citrate Buffer, pH 6.0, for 10-20 min followed by cooling at RT for 20 minutes),




Fig. 1: Formalin-fixed, paraffin-embedded human Tonsil Dendritic stained with CD21-Monospecific Mouse Monoclonal Antibody (CR2/2754).




Fig. 2: SDS-PAGE Analysis Purified CD21-Monospecific Mouse Monoclonal Antibody (CR2/2754). Confirmation of Integrity and Purity of Antibody.




Fig. 3: Western Blot Analysis of human Raji cell lysate using CD21-Monospecific Mouse Monoclonal Antibody (CR2/2754).




Fig. 4: Flow Cytometric Analysis of PFA-fixed MOLT4 cells. CD21-Monospecific Mouse Monoclonal Antibody (CR2/2754) followed by Goat anti-Mouse IgG-CF488 (Blue); Isotype Control (Red).




Fig. 5: Immunofluorescence staining of PFA-fixed MOLT4 cells using CD21-Monospecific Mouse Monoclonal Antibody (CR2/2754) followed by goat anti-Mouse IgG conjugated to CF488 (green). Nuclei are stained with Reddot.




Fig. 6: Analysis of Protein Array containing more than 19,000 full-length human proteins using CD21-Monospecific Mouse Monoclonal Antibody (CR2/2754) Z- and S-Score: The Z-score represents the strength of a signal that a monoclonal antibody (MAb) (in combination with a fluorescently-tagged anti-IgG secondary antibody) produces when binding to a particular protein on the HuProt™ array. Z-scores are described in units of standard deviations (SD's) above the mean value of all signals generated on that array. If targets on HuProt™ are arranged in descending order of the Z-score, the S-score is the difference (also in units of SD's) between the Z-score. S-score therefore represents the relative target specificity of a MAb to its intended target. A MAb is considered to specific to its intended target, if the MAb has an S-score of at least 2.5. For example, if a MAb binds to protein X with a Z-score of 43 and to protein Y with a Z-score of 14, then the S-score for the binding of that MAb to protein X is equal to 29.