

**36-2122: Anti-GM-CSF (Granulocyte/Macrophage - Colony Stimulating Factor) Monoclonal Antibody(Clone: SPM469)**

<b>Clonality :</b>	Monoclonal
<b>Clone Name :</b>	SPM469
<b>Application :</b>	FACS,IF
<b>Reactivity :</b>	Human
<b>Gene :</b>	CSF2
<b>Gene ID :</b>	1437
<b>Uniprot ID :</b>	P04141
<b>Alternative Name :</b>	Burst Promoting Activity; Colony stimulating factor 2 (granulocyte-macrophage); Eosinophil Colony Stimulating Factor; Granulocyte Macrophage Colony Stimulating Factor; Molgramostin; Pluripoietin Alpha; Sargramostim
<b>Isotype :</b>	Rat IgG2a, kappa
<b>Immunogen Information :</b>	Recombinant human GM-CSF protein

**Description**

Granulocyte/macrophage - Colony-stimulating factor (GM-CSF) is a hematopoietic factor that is produced by activated T-cells, B-cells, mast cells, macrophages, fibroblasts, and endothelial cells. In addition to supporting colony formation of granulocyte/macrophage progenitors, GM-CSF is a growth factor for erythroid, megakaryocyte, and eosinophil progenitors.

**Product Info**

<b>Amount :</b>	20 µg / 100 µg
<b>Content :</b>	200 µg/ml of Ab Purified from Bioreactor Concentrate by Protein A/G. Prepared in 10mM PBS with 0.05% BSA & 0.05% azide. Also available WITHOUT BSA & azide at 1.0mg/ml.
<b>Storage condition :</b>	Antibody with azide - store at 2 to 8°C. Antibody without azide - store at -20 to -80°C. Antibody is stable for 24 months. Non-hazardous.

**Application Note**

Neutralization Studies (Order Ab without Azide); Flow Cytometry (1-2ug/million cells); Immunofluorescence (1-2ug/ml);