

9853 Pacific Heights Blvd. Suite D. San Diego, CA 92121, USA Tel: 858-263-4982

Email: info@abeomics.com

## 36-2155: Anti-CD55 / Decay Accelerating Factor (DAF) Monoclonal Antibody(Clone: 143-30)

Clonality: Monoclonal
Clone Name: 143-30
Application: FACS,IF
Reactivity: Human
Gene: CD55
Gene ID: 1604
Uniprot ID: P08174

CD55; CD55 Cromer blood group system; Complement decay-accelerating factor; CROM; DAF;

Alternative Name: GPI-DAF; Decay Accelerating Factor for Complement; Decay accelerating factor GPI-form;

Decay accelerating factor soluble-form

**Isotype:** Mouse IgG1, kappa

Immunogen Information: PHA stimulated human PBL

## **Description**

Recognizes a single chain glycoprotein of 70kDa, identified as CD55/DAF. It is widely expressed on cells throµghout the body including leukocytes, erythrocytes, epithelium, endothelium, and fibroblasts. It is a Glycosyl phosphatidylinositol anchored (GPI-anchored) member of the membrane bound complement regulatory proteins that inhibit autologous complement cascade activation. It prevents the amplification steps of the complement cascade by interfering with the assembly of the C3-convertases, C4b2a and C3bBb, and the C5-convertase, C4b2a3b and C3bBb3b. CD55 also serves as receptor for CD97 and for echovirus and Coxsackie B virus. The MAb 143-30 may be used as marker for paroxysmal nocturnal hemoglobinuria (PNH).

## **Product Info**

**Amount:** 20 μg / 100 μg

Content: 200 µg/ml of Ab Purified from Bioreactor Concentrate by Protein A/G. Prepared in 10mM PBS

with 0.05% BSA & 0.05% azide. Also available WITHOUT BSA & azide at 1.0mg/ml.

**Storage condition :** Antibody with azide - store at 2 to 8°C. Antibody without azide - store at -20 to -80°C. Antibody

is stable for 24 months. Non-hazardous.

## **Application Note**

Flow Cytometry (1-2ug/million cells); ,Immunofluorescence (1-2ug/ml),




Fig. 1: Immunofluorescent staining of paraformaldehyde-fixed HeLa cells with CD55 Mouse Monoclonal Antibody (143-30) followed by goat anti-Mouse IgG-CF488 (Green). Nuclei are stained with RedDot.



9853 Pacific Heights Blvd. Suite D. San Diego, CA 92121, USA Tel: 858-263-4982

Email: info@abeomics.com




Fig. 2: Flow Cytometric Analysis of paraformaldehyde-fixed HeLa cells with CD55 Mouse Monoclonal Antibody (143-30) followed by goat anti-Mouse IgG-CF488 (Blue) Isotype Control (Red)




Fig. 3: SDS-PAGE Analysis Purified CD55 Mouse Monoclonal Antibody (143-30). Confirmation of Integrity and Purity of Antibody.