

36-2156: Anti-Drebrin 1 (DBN1) Monoclonal Antibody(Clone: DBN1/2879)

Clonality :	Monoclonal
Clone Name :	DBN1/2879
Application :	IHC,WB
Reactivity :	Human
Gene :	DBN1
Gene ID :	1627
Uniprot ID :	Q16643
Alternative Name :	Developmentally regulated brain protein; Drebrin 1; Drebrin; Drebrin E; Drebrin E2; Drebrin1; DrebrinE
Isotype :	Mouse IgG1, kappa
Immunogen Information :	Recombinant fragment (aa 150-281) of human DBN1 protein (exact sequence is proprietary)

Description

Drebrins (developmentally regulated brain proteins) are cytoplasmic proteins that bind F-actin in the brain and are involved in cell migration, extension of neuronal processes and plasticity of dendrites. There are three isoforms: two embryonic types (E1 and E2); and an adult type (A), generated by alternative RNA splicing from a single Drebrin gene. Drebrins are expressed mainly in brain neurons but are also found in skeletal muscle, heart, placenta, pancreas and kidney. Drebrin has been designated as a marker of the dendritic spine. Decreases in Drebrin levels in the brain have been associated with Alzheimer's disease and Down syndrome.

Product Info

Amount :	20 µg / 100 µg
Content :	200 µg/ml of Ab Purified from Bioreactor Concentrate. Prepared in 10mM PBS with 0.05% BSA & 0.05% azide. Also available WITHOUT BSA & azide at 1.0mg/ml.
Storage condition :	Antibody with azide - store at 2 to 8°C. Antibody without azide - store at -20 to -80°C. Antibody is stable for 24 months. Non-hazardous.

Application Note

Western Blot (1-2ug/ml); Immunohistochemistry (Formalin-fixed) (1-2ug/ml for 30 min at RT),(Staining of formalin-fixed tissues requires heating tissue sections in 10mM Tris with 1mM EDTA, pH 9.0, for 45 min at 95°C followed by cooling at RT for 20 minutes);




Fig. 1: Formalin-fixed, paraffin-embedded human Brain stained with Drebrin-1 Mouse Monoclonal Antibody (DBN1/2879).




Fig. 2: Formalin-fixed, paraffin-embedded human Cervix stained with Drebrin-1 Mouse Monoclonal Antibody (DBN1/2879).




Fig. 3: Analysis of Protein Array containing more than 19,000 full-length human proteins using Drebrin-1 Mouse Monoclonal Antibody (DBN1/2879). Z- and S- Score: The Z-score represents the strength of a signal that a monoclonal antibody (Monoclonal Antibody) (in combination with a fluorescently-tagged anti-IgG secondary antibody) produces when binding to a particular protein on the HuProt™ array. Z-scores are described in units of standard deviations (SD's) above the mean value of all signals generated on that array. If targets on HuProt™ are arranged in descending order of the Z-score, the S-score is the difference (also in units of SD's) between the Z-score. S-score therefore represents the relative target specificity of a Monoclonal Antibody to its intended target. A Monoclonal Antibody is considered to specific to its intended target, if the Monoclonal Antibody has an S-score of at least 2.5. For example, if a Monoclonal Antibody binds to protein X with a Z-score of 43 and to protein Y with a Z-score of 14, then the S-score for the binding of that Monoclonal Antibody to protein X is equal to 29.