

## 36-2191: Anti-Thymidine Phosphorylase / PD-ECGF (Angiogenesis Marker) Monoclonal Antibody(Clone: TYMP/2890R)

|                                |  |
|--------------------------------|--|
| <b>Clonality :</b>             | Monoclonal   |
| <b>Clone Name :</b>            | TYMP/2890R   |
| <b>Application :</b>           | WB,IHC   |
| <b>Reactivity :</b>            | Human, Mouse, Rat  |
| <b>Gene :</b>                  | TYMP   |
| <b>Gene ID :</b>               | 1890   |
| <b>Uniprot ID :</b>            | P19971   |
| <b>Alternative Name :</b>      | ECGF; ECGF1; Gliostatin; hPD-ECGF; MEDPS1; MNGIE; MTDPS1; PD-ECGF; PDECGF; Platelet-derived endothelial cell growth factor; TdRPase; Thymidine phosphorylase; TP; Tymp |
| <b>Isotype :</b>               | Rabbit IgG   |
| <b>Immunogen Information :</b> | Recombinant full-length human Thymidine Phosphorylase (TP / PD-ECGF) protein   |

### Description

Recognizes a protein (amino acid 482) of 55kDa (in vivo 110kDa homodimer), identified as platelet-derived endothelial growth factor (PD-ECGF), same as thymidine phosphorylase (TP) or gliostatin. In the presence of inorganic orthophosphate, it catalyzes the reversible phospholytic cleavage of thymidine and deoxyuridine to their corresponding bases and 2-deoxyribose-1-phosphate. It is both chemotactic and mitogenic for endothelial cells and a non-heparin binding angiogenic factor present in platelets. Its enzymatic activity is crucial for angiogenic activity (metabolite is angiogenic). Higher levels of serum TP/PD-ECGF are observed in cancer patients. It is also involved in transformation of fluoropyrimidines, cytotoxic agents used in the treatment of a variety of malignancies, into active cytotoxic metabolites (e.g. 5'-deoxy-5-fluorouridine to 5-FU). High intra-cellular levels of TP/PD-ECGF are associated with increased chemosensitivity to such antimetabolites.

### Product Info

|                            |   |
|----------------------------|---|
| <b>Amount :</b>            | 20 µg / 100 µg  |
| <b>Content :</b>           | 200 µg/ml of Ab Purified from Bioreactor Concentrate by Protein A/G. Prepared in 10mM PBS with 0.05% BSA & 0.05% azide. Also available WITHOUT BSA & azide at 1.0mg/ml. |
| <b>Storage condition :</b> | Antibody with azide - store at 2 to 8°C. Antibody without azide - store at -20 to -80°C. Antibody is stable for 24 months. Non-hazardous.                               |

### Application Note

Western Blot (2-4ug/ml); Immunohistochemistry (Formalin-fixed) (1-2ug/ml for 30 minutes at RT)(Staining of formalin-fixed tissues requires heating tissue sections in 10mM Tris with 1mM EDTA, pH 9.0, for 45 min at 95°C followed by cooling at RT for 20 minutes);




Fig. 1: Formalin-fixed, paraffin-embedded human Prostate Carcinoma stained with Thymidine Phosphorylase Rabbit Recombinant Monoclonal (TYMP/2890R).




Fig. 2: SDS-PAGE Analysis of Purified Thymidine Phosphorylase Rabbit Recombinant Monoclonal (TYMP/2890R). Confirmation of Purity and Integrity of Antibody.