

36-2212: Anti-TIGIT / VSTM3 / VSIG9 (Immune Checkpoint for Cancer) Monoclonal Antibody(Clone: TIGIT/3017)

Clonality :	Monoclonal
Clone Name :	TIGIT/3017
Application :	ELISA,FACS,IHC
Reactivity :	Human
Gene :	TIGIT
Gene ID :	201633
Uniprot ID :	Q4951
Alternative Name :	T-cell immunoreceptor with Ig and ITIM domains; V-set and immunoglobulin domain containing 9 (VSIG9); V-set and transmembrane domain containing 3 (VSTM3); Washington University cell adhesion molecule (WUCAM)
Isotype :	Mouse IgG2c, kappa
Immunogen Information :	Recombinant fragment (around aa 22-141) of human TIGIT protein (exact sequence is proprietary)

Description

TIGIT is a checkpoint inhibitor which binds with high affinity to the poliovirus receptor (PVR), causing increased IL10 secretion, decreased IL12B secretion. TIGIT binding to PVR also causes the suppression of T cell activation by promoting the generation of mature immuno-regulatory dendritic cells. It is expressed at low levels on natural killer (NK) cells, as well as peripheral memory and regulatory CD4+ T cells. At the protein level, it is upregulated following the activation of these cells. Functionally, TIGIT is similar to CTLA4. The ligands for TIGIT include CD155 (signal abrogation) and CD226 (signal stimulation). It has been demonstrated to be upregulated on T cells in many cancers and is an immuno-oncology target for therapy.

Product Info

Amount :	20 µg / 100 µg
Content :	200 µg/ml of Ab Purified from Bioreactor Concentrate by Protein A/G. Prepared in 10mM PBS with 0.05% BSA & 0.05% azide. Also available WITHOUT BSA & azide at 1.0mg/ml.
Storage condition :	Antibody with azide - store at 2 to 8°C. Antibody without azide - store at -20 to -80°C. Antibody is stable for 24 months. Non-hazardous.

Application Note

ELISA (For coating, order antibody without BSA); Flow Cytometry (1-2ug/million cells); Immunohistochemistry (Formalin-fixed) (1-2ug/ml for 30 minutes at RT),(Staining of formalin-fixed tissues requires heating tissue sections in 10mM Tris with 1mM EDTA, pH 9.0, for 45 min at 95°C followed by cooling at RT for 20 minutes),




Fig. 1: Formalin-fixed, paraffin-embedded human Prostate Carcinoma stained with TIGIT-Monospecific Mouse Monoclonal Antibody (TIGIT/3017).




Fig. 2: SDS-PAGE Analysis Purified Monospecific Mouse Monoclonal Antibody to TIGIT (TIGIT/3017). Confirmation of Integrity and Purity of Antibody.




Fig. 3: Flow Cytometric Analysis of PFA-fixed MOLT4 cells. TIGIT-Monospecific Mouse Monoclonal Antibody (TIGIT/3017) followed by goat anti-Mouse IgG-CF488 (Blue); Isotype Control (Red).




Fig. 4: Analysis of Protein Array containing more than 19,000 full-length human proteins using TIGIT-Monospecific Mouse Monoclonal Antibody (TIGIT/3017). Z- and S-Score: The Z-score represents the strength of a signal that a monoclonal antibody (MAb) (in combination with a fluorescently-tagged anti-IgG secondary antibody) produces when binding to a particular protein on the HuProt™ array. Z-scores are described in units of standard deviations (SD's) above the mean value of all signals generated on that array. If targets on HuProt™ are arranged in descending order of the Z-score, the S-score is the difference (also in units of SD's) between the Z-score. S-score therefore represents the relative target specificity of a MAb to its intended target. A MAb is considered to specific to its intended target, if the MAb has an S-score of at least 2.5. For example, if a MAb binds to protein X with a Z-score of 43 and to protein Y with a Z-score of 14, then the S-score for the binding of that MAb to protein X is equal to 29.