w abeomics

36-2214: Anti-Endoglin / CD105 (Angiogenesis Marker) Monoclonal Antibody(Clone: ENG/3269)

Clonality :	Monoclonal
Clone Name :	ENG/3269
Application :	IHC
Reactivity :	Human
Gene :	ENG
Gene ID :	2022
Uniprot ID :	P17813
Alternative Name :	CD105; END; Endoglin; Eng; HHT1; S-endoglin
Isotype :	Mouse IgG2b, kappa
Immunogen Information :	Recombinant human Endoglin protein fragment (aa74-251) (exact sequence is proprietary)

Description

Product Info

CD105/Endoglin is a Type I membrane glycoprotein located on cell surfaces and is part of the TGF-beta receptor complex. This protein has been found on endothelial cells, activated macrophages, fibroblasts, and smooth-muscle cells. Endoglin has a role in the development of the cardiovascular system and in vascular remodeling. Its expression is regulated during heart development. CD105 is highly expressed in endothelial cells during tumor angiogenesis and inflammation, with weak or negative expression in vascular endothelium of normal tissues. Angiogenesis is a promising prognostic marker in a variety of tumors. Endoglin is a more specific and sensitive marker for tumor angiogenesis than CD31 or CD34, as it labels only newly-formed blood vessels and may serve as a prognostic marker for Prostate Adenocarcinoma, and cancers of the lung, stomach, breast, and brain.

20 μg / 100 μg
200 μg/ml of Ab Purified from Bioreactor Concentrate by Protein A/G. Prepared in 10mM PBS with 0.05% BSA & 0.05% azide. Also available WITHOUT BSA & azide at 1.0mg/ml.
Antibody with azide - store at 2 to 8°C. Antibody without azide - store at -20 to -80°C. Antibody is stable for 24 months. Non-hazardous.

Application Note

Immunohistochemistry (Formalin-fixed) (1-2ug/ml for 30 minutes at RT),(Staining of formalin-fixed tissues requires heating tissue sections in 10mM Tris with 1mM EDTA, pH 9.0, for 45 min at 95°C followed by cooling at RT for 20 minutes);



Fig. 1: Formalin-fixed, paraffin-embedded human Tonsil stained with Endoglin / CD105 Mouse Monoclonal Antibody (ENG/3269).

For Research Use Only. Not for use in diagnostic/therapeutics procedures.

w abeomics

9853 Pacific Heights Blvd. Suite D. San Diego, CA 92121, USA Tel: 858-263-4982 Email: info@abeomics.com



Fig. 2: Formalin-fixed, paraffin-embedded human Tonsil stained with Endoglin / CD105 Mouse Monoclonal Antibody (ENG/3269).



Fig. 3: SDS-PAGE Analysis Purified Endoglin / CD105 Mouse Monoclonal Antibody (ENG/3269). Confirmation of Integrity and Purity of Antibody.



Fig. 4: Analysis of Protein Array containing more than 19,000 full-length human proteins using Endoglin / CD105 Mouse Monoclonal Antibody (ENG/3269). Z- and S-Score: The Z-score represents the strength of a signal that a monoclonal antibody (MAb) (in combination with a fluorescently-tagged anti-IgG secondary antibody) produces when binding to a particular protein on the HuProtTM array. Z-scores are described in units of standard deviations (SD's) above the mean value of all signals generated on that array. If targets on HuProtTM are arranged in descending order of the Z-score, the S-score is the difference (also in units of SD's) between the Z-score. S-score therefore represents the relative target specificity of a MAb to its intended target. A MAb is considered to specific to its intended target, if the MAb has an Sscore of at least 2.5. For example, if a MAb binds to protein X with a Z-score of 43 and to protein Y with a Z-score of 14, then the S-score for the binding of that MAb to protein X is equal to 29.