

9853 Pacific Heights Blvd. Suite D. San Diego, CA 92121, USA Tel: 858-263-4982

Email: info@abeomics.com

36-2218: Anti-NSE gamma (Neuron Specific Enolase, gamma) (Neuroendocrine Marker) Monoclonal Antibody(Clone: NSE-P2)

Clonality: Monoclonal
Clone Name: NSE-P2
Application: IHC,FACS,IF
Reactivity: Human
Gene: ENO2
Gene ID: 2026
Uniprot ID: P09104

2-phospho-D-glycerate hydrolyase; ENO2; ENOG; Enolase 2 gamma neuronal; Enolase2;

Alternative Name: Gamma-enolase; Neural enolase; Neuron specific gamma enolase; Neuron-specific enolase;

NSE

Isotype: Mouse IgG1, kappa

Immunogen Information: A synthetic peptide of human NSE gamma (around aa416-433) (exact sequence is

proprietary)

Description

This monoclonal antibody recognizes a protein of about 50kDa, which is identified as gamma-enolase. Three isoenzymes of enolases are identified, alpha, beta and gamma. Alpha-isoform is expressed in most tissues, whereas beta-form is expressed predominantly in muscle tissue whereas gamma-enolase is found only in nervous tissue. These isoforms exist as both homodimers and heterodimers, and they play a role in converting phosphoglyceric acid to phosphenolpyruvic acid in the glycolytic pathway. NSE-gamma is a useful marker to identify peripheral nerves and tumors of neuro-endocrine origins, such as pheochromocytomas.It it be usually employed in combination with other markers such as Synaptophysin, Chromogranin A, and Neurofilament.

Product Info

Amount : 20 μg / 100 μg

Content: 200 µg/ml of Ab Purified from Bioreactor Concentrate by Protein A/G. Prepared in 10mM PBS

with 0.05% BSA & 0.05% azide. Also available WITHOUT BSA & azide at 1.0mg/ml.

Storage condition : Antibody with azide - store at 2 to 8°C. Antibody without azide - store at -20 to -80°C. Antibody

is stable for 24 months. Non-hazardous.

Application Note

Flow Cytometry (1-2ug/million cells); Immunofluorescence (1-2ug/ml); Immunohistochemistry (Formalin-fixed) (1-2ug/ml for 30 min at RT)(Staining of formalin-fixed tissues requires heating tissue sections in 10mM Tris with 1mM EDTA, pH 9.0, for 45 min at 95°C followed by cooling at RT for 20 minutes);



9853 Pacific Heights Blvd. Suite D. San Diego, CA 92121, USA Tel: 858-263-4982

Email: info@abeomics.com




Fig. 1: Formalin-fixed, paraffin-embedded human Neuroendocrine tumor stained with NSE gamma Mouse Monoclonal Antibody (NSE-P2).




Fig. 2: SDS-PAGE Analysis Purified NSE gamma Mouse Monoclonal Antibody (NSE-P2). Confirmation of Purity and Integrity of Antibody.