

36-2272: Anti-Factor XIIIa (Coagulation Factor XIIIa Chain) Monoclonal Antibody(Clone: F13A1/3772R)

Clonality :	Monoclonal
Clone Name :	F13A1/3772R
Application :	IHC
Reactivity :	Human
Gene :	F13A1
Gene ID :	2162
Uniprot ID :	P00488
Alternative Name :	Coagulation factor XIII A chain; Coagulation factor XIII A1 polypeptide; F13A; F13a1; Fibrin stabilizing factor, A subunit; Fibrinolyase; FSF, A subunit; Protein-glutamine gamma-glutamyltransferase A chain; TGase; Transglutaminase. plasma
Isotype :	Rabbit IgG
Immunogen Information :	Recombinant fragment of human Factor XIIIa protein (aa46-181) (exact sequence is proprietary)

Description

It recognizes a protein of 83kDa, which is identified as Factor XIIIa. It has been identified in platelets, megakaryocytes, and fibroblast-like mesenchymal or histiocytic cells in the placenta, uterus, and prostate, monocytes and macrophages and dermal dendritic cells. Anti-factor XIIIa has been found to be useful in differentiating between dermatofibroma (almost all cases are positive), dermatofibrosarcoma protuberans (-/+) and desmoplastic malignant melanoma (-). Anti-factor XIIIa positivity is also seen in capillary hemangioblastoma, hemangioendothelioma, hemangiopericytoma, xanthogranuloma, xanthoma, hepatocellular carcinoma, glomus tumor, and meningioma.

Product Info

Amount :	20 µg / 100 µg
Content :	200 µg/ml of Ab Purified from Bioreactor Concentrate by Protein A/G. Prepared in 10mM PBS with 0.05% BSA & 0.05% azide. Also available WITHOUT BSA & azide at 1.0mg/ml.
Storage condition :	Antibody with azide - store at 2 to 8°C. Antibody without azide - store at -20 to -80°C. Antibody is stable for 24 months. Non-hazardous.

Application Note

Immunohistochemistry (Formalin-fixed) (1-2ug/ml for 30 min at RT) ,(Staining of formalin-fixed tissues requires heating tissue sections in 10mM Tris with 1mM EDTA, pH 9.0, for 45 min at 95°C followed by cooling at RT for 20 minutes);




Fig. 1: Formalin-fixed, paraffin-embedded human placenta stained with Factor 13A Recombinant Rabbit Monoclonal Antibody (F13A1/3772R).




Fig. 2: Formalin-fixed, paraffin-embedded human placenta stained with Factor 13A Recombinant Rabbit Monoclonal Antibody (F13A1/3772R).