

9853 Pacific Heights Blvd. Suite D. San Diego, CA 92121, USA Tel: 858-263-4982

Email: info@abeomics.com

36-2272: Anti-Factor XIIIa (Coagulation Factor XIIIA Chain) Monoclonal Antibody(Clone: F13A1/3772R)

Clonality: Monoclonal **Clone Name:** F13A1/3772R

Application: IHC Reactivity: Human Gene: F13A1 Gene ID: 2162 **Uniprot ID:** P00488

Coagulation factor XIII A chain; Coagulation factor XIII A1 polypeptide; F13A; F13a1; Fibrin **Alternative Name:**

stabilizing factor, A subunit; Fibrinoligase; FSF, A subunit; Protein-glutamine gamma-

glutamyltransferase A chain; TGase; Transglutaminase. plasma

Isotype: Rabbit IaG

Recombinant fragment of human Factor XIIIa protein (aa46-181) (exact sequence is Immunogen Information:

proprietary)

Description

It recognizes a protein of 83kDa, which is identified as Factor XIIIa. It has been identified in platelets, megakaryocytes, and fibroblast-like mesenchymal or histiocytic cells in the placenta, uterus, and prostate, monocytes and macrophages and dermal dendritic cells. Anti-factor XIIIa has been found to be useful in differentiating between dermatofibroma (almost all cases are positive), dermatofibrosarcoma protuberans (-/+) and desmoplastic malignant melanoma (-). Anti-factor XIIIa positivity is also seen in capillary hemagioblastoma, hemangioendothelioma, hemangiopericytoma, xanthogranuloma, xanthoma, hepatocellular carcinoma, glomus tumor, and meningioma.

Product Info

Amount: 20 μg / 100 μg

200 µg/ml of Ab Purified from Bioreactor Concentrate by Protein A/G. Prepared in 10mM PBS Content:

with 0.05% BSA & 0.05% azide. Also available WITHOUT BSA & azide at 1.0mg/ml.

Antibody with azide - store at 2 to 8°C. Antibody without azide - store at -20 to -80°C. Antibody Storage condition:

is stable for 24 months. Non-hazardous.

Application Note

Immunohistochemistry (Formalin-fixed) (1-2ug/ml for 30 min at RT) ,(Staining of formalin-fixed tissues requires heating tissue sections in 10mM Tris with 1mM EDTA, pH 9.0, for 45 min at 95&deqC followed by cooling at RT for 20 minutes);




Fig. 1: Formalin-fixed, paraffin-embedded human placenta stained with Factor 13A Recombinant Rabbit Monoclonal Antibody (F13A1/3772R).



9853 Pacific Heights Blvd. Suite D. San Diego, CA 92121, USA Tel: 858-263-4982

Email: info@abeomics.com




Fig. 2: Formalin-fixed, paraffin-embedded human placenta stained with Factor 13A Recombinant Rabbit Monoclonal Antibody (F13A1/3772R).