

## 36-2281: Anti-CD32 (Fc Gamma RIIa) Monoclonal Antibody(Clone: 7.3)

<b>Clonality :</b>	Monoclonal
<b>Clone Name :</b>	7.3
<b>Application :</b>	Functional Assay,IF,FACS
<b>Reactivity :</b>	Human
<b>Gene :</b>	FCGR2A
<b>Gene ID :</b>	2212
<b>Uniprot ID :</b>	P12318
<b>Alternative Name :</b>	CD32A; Fc fragment of IgG low affinity IIa receptor; Fc gamma RIIa; Fc-gamma RII-a; Fc-gamma-RIIa; FCGR2A; FcgRII; Fcr-2; FcRII-a; IGFR2; IgG Fc receptor II-a; Low affinity immunoglobulin gamma Fc region receptor II-a; Ly-17; Ly-m20; LyM-1; Lymphocyte antigen 17
<b>Isotype :</b>	Mouse IgG1, kappa
<b>Immunogen Information :</b>	K562 and FcgRII+L cells

### Description

This MAb reacts with a CD32 (FcγRII) epitope (cluster-4). It displays a stronger reaction with Daudi than with U937 cells. The epitope is located in domain 2 of FcγRIIa. Its Fab'2 fragments block immune complex binding. CD32 (FcRII) is a type 1 transmembrane glycoprotein that mediates several functions including phagocytosis, cytotoxicity, and immunomodulation as well as platelet aggregation. Three genes (A, B, and C) encode CD32 and at least 6 isoforms are generated via alternative mRNA splicing, i.e., IIa1, IIa2, IIb1, IIb2, IIb3 and IIc. Monocytes/macrophages, placental trophoblasts and endothelial cells express all isoforms. In addition, the IIb isoform is expressed by B cells, and the IIa isoform by platelets, granulocytes and, weakly, by B cells. NK cells and neutrophils express isoform IIc. CD32 binds weakly to the Fc region of monomeric IgG but more strongly to IgG aggregates and immune complexes.

### Product Info

<b>Amount :</b>	20 µg / 100 µg
<b>Content :</b>	200 µg/ml of Ab Purified from Bioreactor Concentrate by Protein A/G. Prepared in 10mM PBS with 0.05% BSA & 0.05% azide. Also available WITHOUT BSA & azide at 1.0mg/ml.
<b>Storage condition :</b>	Antibody with azide - store at 2 to 8°C. Antibody without azide - store at -20 to -80°C. Antibody is stable for 24 months. Non-hazardous.

### Application Note

Functional Studies (Order Ab without BSA & Azide)Flow Cytometry (1-2ug/million cells); Immunofluorescence (1-2ug/ml);




Fig. 1: SDS-PAGE Analysis Purified CD32 Mouse Monoclonal Antibody (7.30). Confirmation of Purity and Integrity of Antibody.