

## 36-2313: Anti-CELA3B / ELA3B (Pancreatic Function Marker) Monoclonal Antibody(Clone: CELA3B/2810R)

<b>Clonality :</b>	Monoclonal
<b>Clone Name :</b>	CELA3B/2810R
<b>Application :</b>	IHC
<b>Reactivity :</b>	Human
<b>Gene :</b>	CELA3B
<b>Gene ID :</b>	23436
<b>Uniprot ID :</b>	P08861
<b>Alternative Name :</b>	Chymotrypsin like elastase family member 3B (CELA3B); ELA3B; Elastase IIIB; Protease E
<b>Isotype :</b>	Rabbit IgG
<b>Immunogen Information :</b>	Recombinant human CELA3B protein fragment (around aa 82-238) (exact sequence is proprietary)

### Description

This MAb recognizes a protein of ~30kDa, identified as CELA3B (Chymotrypsin like elastase family member 3B). Elastases form a subfamily of serine proteases that hydrolyze many proteins in addition to elastin. Humans have six elastase genes which encode the structurally similar proteins elastase 1, 2, 2A, 2B, 3A, and 3B. Unlike other elastases, elastase 3B has little elastolytic activity. Like most of the human elastases, elastase 3B is secreted from the pancreas as a zymogen and, like other serine proteases such as trypsin, chymotrypsin and kallikrein; it has a digestive function in the intestine. Elastase 3B preferentially cleaves proteins after alanine residues. Elastase 3B may also function in the intestinal transport and metabolism of cholesterol. Both elastase 3A and elastase 3B have been referred to as protease E and as elastase 1, and excretion of this protein in fecal material is frequently used as a measure of pancreatic function.

### Product Info

<b>Amount :</b>	20 µg / 100 µg
<b>Content :</b>	200 µg/ml of Ab Purified from Bioreactor Concentrate by Protein A/G. Prepared in 10mM PBS with 0.05% BSA & 0.05% azide. Also available WITHOUT BSA & azide at 1.0mg/ml.
<b>Storage condition :</b>	Antibody with azide - store at 2 to 8°C. Antibody without azide - store at -20 to -80°C. Antibody is stable for 24 months. Non-hazardous.

### Application Note

Immunohistochemistry (Formalin-fixed) (1-2ug/ml for 30 minutes at RT)(Staining of formalin-fixed tissues requires heating tissue sections in 10mM Tris with 1mM EDTA, pH 9.0, for 45 min at 95&degC followed by cooling at RT for 20 minutes);




Fig. 1: Formalin-fixed, paraffin-embedded human Pancreas stained with CELA3B Rabbit Recombinant Monoclonal Antibody (CELA3B/2810R).




Fig. 2: SDS-PAGE Analysis of Purified CELA3B Rabbit Recombinant Monoclonal Antibody (CELA3B/2810R). Confirmation of Purity and Integrity of Antibody.