

## 36-2357: Anti-CD15 / FUT4 (Reed-Sternberg Cell Marker) Polyclonal Antibody

<b>Clonality :</b>	Polyclonal
<b>Application :</b>	FACS,IF,IHC
<b>Reactivity :</b>	Human, Mouse, Rat
<b>Gene :</b>	FUT4
<b>Gene ID :</b>	2526
<b>Uniprot ID :</b>	P22083
<b>Alternative Name :</b>	3 Fucosyl N Acetyl Lactosamine; Alpha (1,3) Fucosyltransferase; Alpha 13 fucosyltransferase FucT; ELAM Ligand Fucosyltransferase; ELFT; FCT3A; Fuc-TIV; Fucosyltransferase 4 Alpha 1 3 Fucosyltransferase Myeloid Specific; Fucosyltransferase 4; Galactoside 3 L Fucosyltransferase; Lewis X; LeX; SSEA1; Stage Specific Embryonic Antigen 1
<b>Isotype :</b>	Rabbit IgG
<b>Immunogen Information :</b>	Purified human neutrophils

### Description

CD15 plays a role in mediating phagocytosis, bactericidal activity, and chemotaxis. It is present on s disease. CD15 is occasionally expressed in large cell lymphomas of both B and T phenotypes which otherwise have a quite distinct histological appearance.

### Product Info

<b>Amount :</b>	20 µg / 100 µg
<b>Content :</b>	200 µg/ml of Ab Purified by Protein A Column. Prepared in 10mM PBS with 0.05% BSA & 0.05% azide. Also available WITHOUT BSA & Azide at 1.0mg/ml.
<b>Storage condition :</b>	Antibody with azide - store at 2 to 8°C. Antibody without azide - store at -20 to -80°C. Antibody is stable for 24 months. Non-hazardous.

### Application Note

Flow Cytometry (2-4ug/million cells); Immunofluorescence (2-4ug/ml); Immunohistochemistry (Formalin-fixed) (5-10ug/ml for 30 minutes at RT)(Staining of formalin-fixed tissues requires heating tissue sections in 10mM Tris with 1mM EDTA, pH 9.0, for 45 min at 95&degC followed by cooling at RT for 20 minutes)




Fig. 1: Formalin-fixed, paraffin-embedded human Hodgkin's Lymphoma stained with CD15 Rabbit Polyclonal Antibody.