

36-2390: Anti-Glypican-3 (GPC3) (Hepatocellular Carcinoma Marker) Monoclonal Antibody(Clone: SPM595)-CF555

| | |
|--------------------------------|---|
| Clonality : | Monoclonal |
| Clone Name : | SPM595 |
| Application : | FACS,IF,IHC |
| Reactivity : | Human |
| Conjugate : | CF555 |
| Gene : | GPC3 |
| Gene ID : | 2719 |
| Uniprot ID : | P51654 |
| Alternative Name : | DGSX; Glypican proteoglycan 3; GPC3; GTR2-2; Heparan sulphate proteoglycan; Intestinal protein OCI-5; MXR7; OCI-5; SDYS; Secreted glypican-3; SGBS1 |
| Isotype : | Mouse IgG1, kappa |
| Immunogen Information : | A recombinant fragment containing amino acids 511-580 of human glypican-3 |

Description

Glypican-3 (GPC3) is a glycosylphosphatidyl inositol-anchored membrane protein, which may also be found in a secreted form. Anti-GPC3 has been identified as a useful tumor marker for the diagnosis of hepatocellular carcinoma (HCC), hepatoblastoma, melanoma, testicular germ cell tumors, and Wilms tumor and hepatoblastoma, with a low or undetectable expression in normal adjacent tissue. In patients with thyroid cancer, expression of GPC3 is dramatically enhanced in certain types of cancers: 100% in follicular carcinoma and 70% in papillary carcinoma. Expression of GPC3 in follicular carcinoma is significantly higher than that of follicular adenoma. In contrast, GPC3 is not expressed in anaplastic carcinoma.

Product Info

| | |
|----------------------------|---|
| Amount : | 0.5 ml at 100µg/ml |
| Content : | 200 µg/ml of Ab Purified from Bioreactor Concentrate by Protein A/G. Prepared in 10mM PBS with 0.05% BSA & 0.05% azide. Also available WITHOUT BSA & azide at 1.0mg/ml. |
| Storage condition : | Antibody with azide - store at 4 to 8°C. Antibody is stable for 24 months. Non-hazardous. |

Application Note

Flow Cytometry (1-2ug/million cells); Immunofluorescence (1-2ug/ml); Immunohistochemistry (Formalin-fixed) (1-2ug/ml for 30 minutes at RT)(Staining of formalin-fixed tissues requires heating tissue sections in 10mM Tris with 1mM EDTA, pH 9.0, for 45 min at 95°C followed by cooling at RT for 20 minutes);