

**36-2418: Anti-CD13 / Aminopeptidase-N (Myeloid Cell Marker) Monoclonal Antibody(Clone: B-F10)**

<b>Clonality :</b>	Monoclonal
<b>Clone Name :</b>	B-F10
<b>Application :</b>	FACS,IF,IHC
<b>Reactivity :</b>	Human
<b>Gene :</b>	ANPEP
<b>Gene ID :</b>	290
<b>Uniprot ID :</b>	P15144
<b>Alternative Name :</b>	Alanyl aminopeptidase; Aminopeptidase M; Aminopeptidase N; ANPEP; AP-M; APN; gp150; hAPN; Lap1; Microsomal aminopeptidase; Myeloid plasma membrane glycoprotein CD13; p150; PEPN
<b>Isotype :</b>	Mouse IgG1, kappa
<b>Immunogen Information :</b>	KG-1 myeloid cells

**Description**

Recognizes an integral membrane glycoprotein of 150kDa, identified as CD13 (also known as aminopeptidase-N). The CD13 antigen is present on most cells of myeloid origin including granulocytes, monocytes, mast cells, and GM-progenitor cells. It is also expressed by the majority of AML, CML in myeloid blast crisis, and in a smaller fraction of lymphoid leukemias. CD13 is absent from normal lymphocytes, platelets and erythrocytes. CD13 is also present on fibroblasts; endothelial cells, epithelial cells from renal proximal tubules and intestinal brush border, bone marrow stromal cells, osteoclasts, and cells lining bile duct canaliculi. CD13 is identical to aminopeptidase N (APN), a prominent membrane-bound metalloprotease present on the surface of intestinal brush border and renal tubules. CD13 plays a role in metabolism of biologically active peptides, in phagocytosis, and in bactericidal/tumoricidal activities. It also serves as a receptor for human coronaviruses (HCV). The lineage-restricted pattern of expression of CD13 within the hemopoietic compartment suggests that it may be important in myeloid cell differentiation.

**Product Info**

<b>Amount :</b>	20 µg / 100 µg
<b>Content :</b>	200 µg/ml of Ab Purified from Bioreactor Concentrate by Protein A/G. Prepared in 10mM PBS with 0.05% BSA & 0.05% azide. Also available WITHOUT BSA & azide at 1.0mg/ml.
<b>Storage condition :</b>	Antibody with azide - store at 2 to 8°C. Antibody without azide - store at -20 to -80°C. Antibody is stable for 24 months. Non-hazardous.

**Application Note**

Flow Cytometry (1-2ug/million cells); Immunofluorescence (1-2ug/ml); Immunohistochemistry (Frozen) (1-2ug/ml for 30 min at RT)