

36-2449: Anti-Granzyme B (NK/T-Cell Lymphoma Marker) Monoclonal Antibody(Clone: GZMB/2403)

Clonality :	Monoclonal
Clone Name :	GZMB/2403
Application :	ELISA
Reactivity :	Human
Gene :	GZMB
Gene ID :	3002
Uniprot ID :	P10144
Alternative Name :	Cathepsin G-like 1; CCPI; CGL1; CSPB; CTLA-1; CTSG1; Cytotoxic serine protease B; Cytotoxic T lymphocyte associated serine esterase 1; Cytotoxic T-lymphocyte proteinase 2; Fragmentin-2; Human lymphocyte protein (Hlp); Lymphocyte protease; SECT; T-cell serine protease 1-3E
Isotype :	Mouse IgG2b, lambda
Immunogen Information :	Recombinant fragment of human GZMB protein (around aa 73-187) (exact sequence is proprietary)

Description

Granzyme B is a member of the granule serine protease family stored specifically in NK cells or cytotoxic T cells. Cytolytic T lymphocytes (CTL) and natural killer (NK) cells share the ability to recognize, bind, and lyse specific target cells. They are thought to protect their host by lysing cells bearing on their surface 'nonself' antigens, usually peptides or proteins resulting from infection by intracellular pathogens. Granzyme B is crucial for the rapid induction of target cell apoptosis by CTLs in the cell-mediated immune response. Granzyme B is useful as a marker in the identification of NK/T-cell lymphomas. High percentages of cytotoxic T-cells have been shown to be an unfavorable prognostic indicator in Hodgkin's Disease.

Product Info

Amount :	20 µg / 100 µg
Content :	200 µg/ml of Ab Purified from Bioreactor Concentrate by Protein A/G. Prepared in 10mM PBS with 0.05% BSA & 0.05% azide. Also available WITHOUT BSA & azide at 1.0mg/ml.
Storage condition :	Antibody with azide - store at 2 to 8°C. Antibody without azide - store at -20 to -80°C. Antibody is stable for 24 months. Non-hazardous.

Application Note

ELISA (For coating use Ab at 1-5ug/ml, order Ab without BSA)




Fig. 1: SDS-PAGE Analysis Purified Granzyme B Mouse Monoclonal Antibody (GZMB/2403). Confirmation of Integrity and Purity of Antibody.




Fig. 2: Analysis of Protein Array containing >19,000 full-length human proteins using Granzyme B Mouse Monoclonal Antibody (GZMB/2403) Z- and S- Score: The Z-score represents the strength of a signal that a monoclonal antibody (MAb) (in combination with a fluorescently-tagged anti-IgG secondary antibody) produces when binding to a particular protein on the HuProtTM array. Z-scores are described in units of standard deviations (SD's) above the mean value of all signals generated on that array. If targets on HuProtTM are arranged in descending order of the Z-score, the S-score is the difference (also in units of SD's) between the Z-score. S-score therefore represents the relative target specificity of a MAb to its intended target. A MAb is considered to specific to its intended target, if the MAb has an S-score of at least 2.5. For example, if a MAb binds to protein X with a Z-score of 43 and to protein Y with a Z-score of 14, then the S-score for the binding of that MAb to protein X is equal to 29.