

## 36-2469: Anti-Heregulin-1 / Neuregulin-1 (Breast and Urothelial Marker) Monoclonal Antibody(Clone: NRG1/2752)

<b>Clonality :</b>	Monoclonal
<b>Clone Name :</b>	NRG1/2752
<b>Application :</b>	ELISA,IF
<b>Reactivity :</b>	Human
<b>Gene :</b>	NRG1
<b>Gene ID :</b>	3084
<b>Uniprot ID :</b>	Q02297
<b>Alternative Name :</b>	Acetylcholine receptor-inducing activity (ARIA); Breast cancer cell differentiation factor p45; Glial growth factor (GGF2); Heregulin-1; Heregulin, alpha (45kD, ERBB2 p185-activator); HGL; HRG1; HRGA; MST131; Neu Differentiation Factor (NDF); Neuregulin-1; NRG1; Sensory and Motor Neuron-Derived Factor (SMDF)
<b>Isotype :</b>	Mouse IgG
<b>Immunogen Information :</b>	Recombinant fragment of human Neuregulin-1 (NRG1) protein (around aa 21-242) (exact sequence is proprietary)

### Description

Heregulin-1 is a membrane glycoprotein that mediates cell-cell signaling and plays a critical role in the growth and development of multiple organ systems. An extraordinary variety of different isoforms are produced from this gene through alternative promoter usage and splicing. These isoforms are expressed in a tissue-specific manner and differ significantly in their structure, and are classified as types I, II, III, IV, V and VI. Dysregulation of this gene has been linked to diseases such as cancer, schizophrenia, and bipolar disorder (BPD).

### Product Info

<b>Amount :</b>	20 µg / 100 µg
<b>Content :</b>	200 µg/ml of Ab Purified from Bioreactor Concentrate by Protein A/G. Prepared in 10mM PBS with 0.05% BSA & 0.05% azide. Also available WITHOUT BSA & azide at 1.0mg/ml.
<b>Storage condition :</b>	Antibody with azide - store at 2 to 8°C. Antibody without azide - store at -20 to -80°C. Antibody is stable for 24 months. Non-hazardous.

### Application Note

ELISA (For coating, order Ab without BSA); Immunofluorescence (1-2ug/ml);




Fig. 1: SDS-PAGE Analysis Purified Heregulin-1 Mouse Monoclonal Antibody (NRG1/2752). Confirmation of Purity and Integrity of Antibody.




Fig. 2: Confocal immunofluorescence of PFA-fixed MCF-7 cells using Heregulin-1 Mouse Monoclonal Antibody (NRG1/2752) followed by Goat-anti Mouse CF488 (green) and Reddot is used to label the nuclei red.




Fig. 3: Analysis of Protein Array containing more than 19,000 full-length human proteins using Heregulin-1 Mouse Monoclonal Antibody (NRG1/2752). Z- and S-Score: The Z-score represents the strength of a signal that a monoclonal antibody (Monoclonal Antibody) (in combination with a fluorescently-tagged anti-IgG secondary antibody) produces when binding to a particular protein on the HuProt™ array. Z-scores are described in units of standard deviations (SD's) above the mean value of all signals generated on that array. If targets on HuProt™ are arranged in descending order of the Z-score, the S-score is the difference (also in units of SD's) between the Z-score. S-score therefore represents the relative target specificity of a Monoclonal Antibody to its intended target. A Monoclonal Antibody is considered to specific to its intended target, if the Monoclonal Antibody has an S-score of at least 2.5. For example, if a Monoclonal Antibody binds to protein X with a Z-score of 43 and to protein Y with a Z-score of 14, then the S-score for the binding of that Monoclonal Antibody to protein X is equal to 29.