

9853 Pacific Heights Blvd. Suite D. San Diego, CA 92121, USA Tel: 858-263-4982

Email: info@abeomics.com

## 36-2479: Anti-HLA-DP (MHC II) Monoclonal Antibody(Clone: SPM421)

Clonality: Monoclonal
Clone Name: SPM421
Application: ELISA,FACS,IF
Reactivity: Human
Gene: HLA-DPB1
Gene ID: 3115

Uniprot ID: P04440

HLA class II histocompatibility antigen DP alpha 1 chain; HLA DP1A; HLA SB alpha chain; HLADP; HLADPA1; HLASB; HLASB histocompatibility type; Human MHC class II HLA SB alpha; HLADPA1; HLADPA1; HLASB histocompatibility type; Human MHC class II HLA SB alpha; HLADPA1; HLADPA1; HLASB histocompatibility type; Human MHC class II HLA SB alpha; HLADPA1; HLADPA1; HLASB histocompatibility type; Human MHC class II HLA SB alpha; HLADPA1; HLADPA1; HLASB histocompatibility type; Human MHC class II HLA SB alpha; HLADPA1; HLADPA1; HLASB histocompatibility type; Human MHC class III histocompatibility type; Human MHC

Major histocompatibility complex class II DP alpha 1; MHC class II DP3 alpha; MHC class II

DPA1; PLT1; Primed lymphocyte test 1

**Isotype:** Mouse IgG2b, kappa

Immunogen Information: Non-T, non-B human acute lymphoblastic leukemia REH6 cell line

## **Description**

Recognizes a non-polymorphic determinant of DP-MHC class II. MHC class II antigens are transmembrane glycoproteins of non-covalently linked alpha (33-35kDa) and beta (27-30kDa) chains. It reportedly reacts with B- non-T, non-B cell lines but not with T- and myeloid cell lines and leukemias. Differential expression of MHC class II antigens on fetal and adult lymphocytes, malignant B cells appears to reflect the stage of cell differentiation which may be useful in the study of lymphoproliferative disorders.

## **Product Info**

**Amount:** 20 μg / 100 μg

**Content:** 200 μg/ml of Ab Purified from Bioreactor Concentrate by Protein A/G. Prepared in 10mM PBS

with 0.05% BSA & 0.05% azide. Also available WITHOUT BSA & azide at 1.0mg/ml.

**Storage condition :** Antibody with azide - store at 2 to 8°C. Antibody without azide - store at -20 to -80°C. Antibody

is stable for 24 months. Non-hazardous.

## **Application Note**

ELISA (For coating, order antibody without BSA); Flow Cytometry (1-2ug/million cells); Immunofluorescence (1-2ug/ml);




Fig. 1: Flow Cytometric Analysis of Human Raji cells using HLA-DP (MHC II) Monoclonal Antibody (SPM421) followed by Goat anti-Mouse IgG-CF488 (Blue); Isotype control (Red).