w abeomics

36-2515: Anti-CD54 / ICAM-1 Monoclonal Antibody(Clone: 15.2)

Clonality :	Monoclonal
Clone Name :	15.2
Application :	Functional Assay, FACS, IF, IHC
Reactivity :	Human
Gene :	ICAM1
Gene ID :	3383
Uniprot ID :	P05362
Alternative Name :	CD54; Cell surface glycoprotein P3.58; Human rhinovirus receptor; ICAM-1; Intercellular adhesion molecule 1; Ly 47; Major group rhinovirus receptor; MALA2; MyD10; Surface antigen of activated B cells
Isotype :	Mouse IgG1, kappa
Immunogen Information : Human monocytes.	

Description

Product Info

Recognizes an 85-115kDa protein (variation with cell type), identified as intercellular adhesion molecule (ICAM-1). It has 7 potential N-linked glycosylation sites. ICAM-1 is a single chain glycoprotein of Ig supergene family, present on unstimulated endothelial cells (EC) and on a variety of other cell types including activated fibroblasts, EC, macrophages, and lymphocytes. ICAM-1 mediates cell adhesion by binding to integrins CD11a/CD18 (leukocyte adhesion molecule, LFA-1) and to CD11b/CD18 (Mac-1). This interaction enhances antigen-specific T-cell activation. ICAM-1 also binds to CD43 and to Plasmodium falciparum infected RBCs. ICAM-1 may also be related to progression and metastasis of tumors.

Amount :20 μg / 100 μgContent :200 μg/ml of Ab Purified from Bioreactor Concentrate by Protein A/G. Prepared in 10mM PBS
with 0.05% BSA & 0.05% azide. Also available WITHOUT BSA & azide at 1.0mg/ml.Storage condition :Antibody with azide - store at 2 to 8°C. Antibody without azide - store at -20 to -80°C. Antibody
is stable for 24 months. Non-hazardous.

Application Note

Functional Studies (Order Ab without BSA & Azide);Flow Cytometry (1-2ug/million cells); Immunofluorescence (1-2ug/ml); Immunohistochemistry (Formalin-fixed) (1-2ug/ml for 30 minutes at RT)(Staining of formalin-fixed tissues requires heating tissue sections in 10mM Tris with 1mM EDTA, pH 9.0, for 45 min at 95°C followed by cooling at RT for 20 minutes);




Fig. 1: SDS-PAGE Analyis Purified CD54 Mouse Monoclonal Antibody (15.2). Confirmation of Purity and Integrity of Antibody.

