

## 36-2537: Anti-IgA (Immunoglobulin Alpha Heavy Chain) (B-Cell Marker) Monoclonal Antibody(Clone: IA761)-CF488

<b>Clonality :</b>	Monoclonal
<b>Clone Name :</b>	IA761
<b>Application :</b>	FACS,IF,IHC
<b>Reactivity :</b>	Human
<b>Conjugate :</b>	CF488
<b>Gene :</b>	IGHA1 & IGHA2
<b>Gene ID :</b>	3493; 3494
<b>Uniprot ID :</b>	P01876; P01877
<b>Alternative Name :</b>	A2m Marker; Ig alpha 1 Chain C Region; Ig alpha 2 Chain C Region; IGHA1; IGHA2; Immunoglobulin Am1; Immunoglobulin Am2; Immunoglobulin Heavy Constant Alpha 1; Immunoglobulin Heavy Constant Alpha 2
<b>Isotype :</b>	Mouse IgG1, kappa
<b>Immunogen Information :</b>	Purified human alpha heavy chain

### Description

This MAbs is specific to heavy chain of IgA and shows minimal cross-reaction with heavy chains of other immunoglobulins. It is reactive with all subclasses of Alpha heavy chain. Immunoglobulins are four-chain, Y-shaped, monomeric structures comprised of two identical heavy chains and two identical light chains held together through inter-chain disulfide bonds. The chains form two domains, the Fab (antigen binding) fragment and the Fc (constant) fragment. Immunoglobulin A (IgA) is the main protein of the mucosal immune system. It is generated by B-cells in gut-associated lymphoid tissues. Daily production of IgA exceeds that of any of the other immunoglobulins. IgA exists mainly in dimers but can also exist as polymers or as monomers. Dimers and polymers contain a joining (J) chain that can be bound by the polymeric immunoglobulin receptor (pIgR) for transportation of the molecule to mucosal surfaces. The most common feature of plasmacytomas, and certain non-Hodgkin's lymphomas is the restricted expression of a single heavy chain class. Demonstration of clonality in lymphoid infiltrates indicates that the infiltrate is clonal and therefore malignant.

### Product Info

<b>Amount :</b>	0.5 ml at 100µg/ml
<b>Content :</b>	200 µg/ml of Ab Purified from Bioreactor Concentrate by Protein A/G. Prepared in 10mM PBS with 0.05% BSA & 0.05% azide. Also available WITHOUT BSA & azide at 1.0mg/ml.
<b>Storage condition :</b>	Antibody with azide - store at 4 to 8°C. Antibody is stable for 24 months. Non-hazardous.

### Application Note

Flow Cytometry (1-2ul/million cells); Immunofluorescence (1-2ug/ml 30 min RT); Immunohistochemistry (Formalin-fixed) (1-2ug/ml 30 min RT) (Staining of formalin-fixed tissues requires heating tissue sections in 10mM Tris with 1mM EDTA, pH 9.0, for 45 min at 95&degC followed by cooling at RT for 20 minutes);




Fig. 1: SDS-PAGE Analysis Purified IgA Mouse Monoclonal Antibody (IA761). Confirmation of Purity and Integrity of Antibody.