w abeomics

36-2606: Anti-CD11b / MAC-1 (Microglial Marker) Monoclonal Antibody(Clone: ITGAM/271)

Clonality :	Monoclonal
Clone Name :	ITGAM/271
Application :	FACS,IF
Reactivity :	Human
Gene :	ITGAM
Gene ID :	3684
Uniprot ID :	P11215
Alternative Name :	CD11 antigen-like family member B; CD11b/CD18; CD49d; Cell surface glycoprotein MAC-1 subunit alpha; Complement Component Receptor 3 Alpha; CR3 Alpha Chain (CR3A); Integrin alpha-M; Integrin beta 2 alpha subunit; ITGAM; Leukocyte adhesion receptor MO1; Ly-40; Mac-1a; MAC1; Mac1, alpha subunit; MAC1A; Macrophage antigen alpha polypeptide; MO1A; Neutrophil adherence receptor alpha M subunit
lsotype :	Mouse IgG1, kappa
Immunogen Information : Recombinant full-length human ITGAM protein	

Description

CD11b is a cell adhesion molecule that acts as a receptor for cell surface ligands such as intracellular adhesion molecules (ICAMs) or soluble ligands. Integrins are heterodimeric proteins that contain an a chain and b chain. Integrin aM combines with the Integrin '2 to form a leukocyte-specific integrin referred to as macrophage receptor 1 (Mac-1), or inactivated-C3b (iC3b) receptor 3 (CR3). Integrin aM/'2 is important in the adherence of neutrophils and monocytes to stimulated endothelium, and also in the phagocytosis of complement coated particles. The protein CD11b has been implicated in the various adhesion-related interactions of cells such as monocytes, macrophages, natural killer (NK) cells, and granulocytes. It is part of a heterodimer that consists of CD11b andCD18. It also modulates the uptake of complement-coated particles within the cell. It is commonly used as a microglial marker in tissues derived from the nervous system.

Product Info

Amount :	20 μg / 100 μg
Content :	200 μg/ml of Ab Purified from Bioreactor Concentrate by Protein A/G. Prepared in 10mM PBS with 0.05% BSA & 0.05% azide. Also available WITHOUT BSA & azide at 1.0mg/ml.
Storage condition :	Antibody with azide - store at 2 to 8°C. Antibody without azide - store at -20 to -80°C. Antibody is stable for 24 months. Non-hazardous.

Application Note

Flow Cytometry (1-2ug/million cells); Immunofluorescence (1-2ug/ml);



9853 Pacific Heights Blvd. Suite D. San Diego, CA 92121, USA Tel: 858-263-4982 Email: info@abeomics.com




Fig. 1: SDS-PAGE Analysis Purified CD11b Mouse Monoclonal Antibody (ITGAM/271). Confirmation of Integrity and Purity of Antibody.