

## 36-2610: Anti-CD11b / MAC-1 (Microglial Marker) Monoclonal Antibody(Clone: M1/70)

<b>Clonality :</b>	Monoclonal
<b>Clone Name :</b>	M1/70
<b>Application :</b>	FACS,IF,Functional Assay
<b>Reactivity :</b>	Human, Mouse
<b>Gene :</b>	ITGAM
<b>Gene ID :</b>	3684
<b>Uniprot ID :</b>	P11215
<b>Alternative Name :</b>	CD11 antigen-like family member B; CD11b/CD18; CD49d; Cell surface glycoprotein MAC-1 subunit alpha; Complement Component Receptor 3 Alpha; CR3 Alpha Chain (CR3A); Integrin alpha-M; Integrin beta 2 alpha subunit; ITGAM; Leukocyte adhesion receptor MO1; Ly-40; Mac-1a; MAC1; Mac1, alpha subunit; MAC1A; Macrophage antigen alpha polypeptide; MO1A; Neutrophil adherence receptor alpha M subunit
<b>Isotype :</b>	Rat IgG2b, kappa
<b>Immunogen Information :</b>	B10 Mouse spleen cells enriched for T lymphocytes

### Description

CD11b is a cell adhesion molecule that acts as a receptor for cell surface ligands such as intracellular adhesion molecules (ICAMs) or soluble ligands. Integrins are heterodimeric proteins that contain an a chain and b chain. Integrin aM combines with the Integrin 2 is important in the adherence of neutrophils and monocytes to stimulated endothelium, and also in the phagocytosis of complement coated particles. The protein CD11b has been implicated in the various adhesion-related interactions of cells such as monocytes, macrophages, natural killer (NK) cells, and granulocytes. It is part of a heterodimer that consists of CD11b and CD18. It also modulates the uptake of complement-coated particles within the cell. It is commonly used as a microglial marker in tissues derived from the nervous system.

### Product Info

<b>Amount :</b>	20 µg / 100 µg
<b>Content :</b>	200 µg/ml of Ab Purified from Bioreactor Concentrate by Protein A/G. Prepared in 10mM PBS with 0.05% BSA & 0.05% azide. Also available WITHOUT BSA & azide at 1.0mg/ml.
<b>Storage condition :</b>	Antibody with azide - store at 2 to 8°C. Antibody without azide - store at -20 to -80°C. Antibody is stable for 24 months. Non-hazardous.

### Application Note

Flow Cytometry (1-2ug/million cells); Immunofluorescence (1-2ug/ml); Functional Studies (Order Ab without BSA & Azide);