

## 36-2631: Anti-Catenin, gamma (Cardiomyocyte Marker) Monoclonal Antibody(Clone: CTNG/1483)

<b>Clonality :</b>	Monoclonal
<b>Clone Name :</b>	CTNG/1483
<b>Application :</b>	IHC
<b>Reactivity :</b>	Human
<b>Gene :</b>	JUP
<b>Gene ID :</b>	3728
<b>Uniprot ID :</b>	P14923
<b>Alternative Name :</b>	ARVD12; Catenin (cadherin-associated protein), gamma 80kDa; Catenin gamma; CTNNG; Desmoplakin III; Desmoplakin-3; DP3; DP111; Junction Plakoglobin; PDGB; PKGB
<b>Isotype :</b>	Mouse IgG1, kappa
<b>Immunogen Information :</b>	Recombinant human gamma catenin protein

### Description

It recognizes a protein of 80-87kDa, identified as gamma-catenin. The catenins (-T-catenin, is located on chromosome 10, and mutations in this gene show a strong correlation to late-onset Alzheimer's disease (LOAD) as well as to dilated cardiomyopathy.

### Product Info

<b>Amount :</b>	20 µg / 100 µg
<b>Content :</b>	200 µg/ml of Ab Purified from Bioreactor Concentrate by Protein A/G. Prepared in 10mM PBS with 0.05% BSA & 0.05% azide. Also available WITHOUT BSA & azide at 1.0mg/ml.
<b>Storage condition :</b>	Antibody with azide - store at 2 to 8°C. Antibody without azide - store at -20 to -80°C. Antibody is stable for 24 months. Non-hazardous.

### Application Note

Immunohistochemistry (Formalin-fixed) (0.1-0.2µg/ml for 30 min at RT)(Staining of formalin-fixed tissues requires heating tissue sections in 10mM Tris with 1mM EDTA, pH 9.0, for 45 min at 95&degC followed by cooling at RT for 20 minutes); with 1mM EDTA, pH 9.0, for 45 min at 95&degC followed by cooling at RT for 20 minutes);




Fig. 1: Formalin-fixed, paraffin-embedded human Cervical Carcinoma stained with Catenin, gamma Monoclonal Antibody (CTNG/1483)




Fig. 2: Formalin-fixed, paraffin-embedded human Skin stained with Catenin, gamma Monoclonal Antibody (CTNG/1483)