

## 36-2639: Anti-PLK1 (Marker of Mitosis) Monoclonal Antibody(Clone: AZ44)

<b>Clonality :</b>	Monoclonal
<b>Clone Name :</b>	AZ44
<b>Application :</b>	ELISA,WB
<b>Gene :</b>	plk1.S
<b>Gene ID :</b>	380481
<b>Uniprot ID :</b>	P70032
<b>Alternative Name :</b>	Cell cycle regulated protein kinase; Polo like kinase 1 (PLK1); Serine/threonine-protein kinase PLK1; STPK13
<b>Isotype :</b>	Mouse IgG1, kappa
<b>Immunogen Information :</b>	Recombinant full-length PLK1 Xenopus laevis protein

### Description

Plk (for polo-like kinase) encodes a serine/threonine kinase that is closely related to polo and CDC5, genes that are required for passage through mitosis in *Drosophila* and *Saccharomyces*, respectively. Polo and Cdc5 both code for proteins that are involved in regulating the function of the mitotic spindle. Plk protein accumulates in the cell during the S and G2 phases of the cell cycle; Plk protein content and catalytic activity peak at the onset of mitosis, followed by a rapid reduction after mitosis. Plk expression is detectable in mitotically active tissues such as colon and placenta, as well as in tumors of various origins. It has also been suggested that Plk may serve as a marker of cell proliferation. Required for recovery after DNA damage checkpoint and entry into mitosis. Required for kinetochore localization of BUB1B. Phosphorylates SGOL1. Required for spindle pole localization of isoform 3 of SGOL1 and plays a role in regulating its centriole cohesion function. Phosphorylates BORA, and thereby promotes the degradation of BORA. Contributes to the regulation of AURKA function. Regulates TP53 stability through phosphorylation of TOPORS. Useful for confirming that extracts have maintained CSF status and have not yet been activated into interphase.

### Product Info

<b>Amount :</b>	20 µg / 100 µg
<b>Content :</b>	200 µg/ml of Ab Purified from Bioreactor Concentrate by Protein A/G. Prepared in 10mM PBS with 0.05% BSA & 0.05% azide. Also available WITHOUT BSA & azide at 1.0mg/ml.
<b>Storage condition :</b>	Antibody with azide - store at 2 to 8°C. Antibody without azide - store at -20 to -80°C. Antibody is stable for 24 months. Non-hazardous.

### Application Note

ELISA (For coating, order antibody without BSA); Western Blot (1-2ug/ml);




Fig. 1: SDS-PAGE Analysis Purified PLK1 Mouse Monoclonal Antibody (AZ44). Confirmation of Purity and Integrity of Antibody.