

36-2672: Anti-Cytokeratin 13 (Non-Keratinized Squamous Epithelial Marker) Monoclonal Antibody(Clone: KRT13/2213)

Clonality :	Monoclonal
Clone Name :	KRT13/2213
Application :	IF,FACS,IHC
Reactivity :	Human
Gene :	KRT13
Gene ID :	3860
Uniprot ID :	P13646
Alternative Name :	CK13; Cytokeratin-13; Keratin Type I Cytoskeletal 13; Keratin-13; KRT13; Type I Cytoskeletal 13; WSN2
Isotype :	Mouse IgG1, kappa
Immunogen Information :	Recombinant full-length human KRT13 protein

Description

Cytokeratin 13 (KRT13) is the major acidic keratin, which together with KRT4, its basic partner, is expressed in the suprabasal layers of non-cornified stratified epithelia including tongue mucosa, esophagus, anal canal epithelium, tracheal epithelium, uterine cervix, and urothelium. Defects in the KRT13 gene are a cause of white sponge nevus of cannon (WSN), a rare autosomal dominant disorder, which predominantly affects non-cornified stratified squamous epithelia and is characterized by the presence of soft, white and spongy plaques in the oral mucosa. KRT13 has been used as a marker for non-keratinized squamous epithelium. It is also expressed in various squamous metaplasia, but it is down regulated in squamous dysplasia and squamous carcinoma.

Product Info

Amount :	20 µg / 100 µg
Content :	200 µg/ml of Ab Purified from Bioreactor Concentrate by Protein A/G. Prepared in 10mM PBS with 0.05% BSA & 0.05% azide. Also available WITHOUT BSA & azide at 1.0mg/ml.
Storage condition :	Antibody with azide - store at 2 to 8°C. Antibody without azide - store at -20 to -80°C. Antibody is stable for 24 months. Non-hazardous.

Application Note

Immunofluorescence (1-2ug/ml); Flow Cytometry (1-2ug per million cells); Immunohistochemistry (Formalin-fixed) (1-2ug/ml for 30 min at RT)(Staining of formalin-fixed tissues requires heating tissue sections in 10mM Tris with 1mM EDTA, pH 9.0, for 45 min at 95°C followed by cooling at RT for 20 minutes);




Fig. 1: Formalin-fixed, paraffin-embedded human Tonsil stained with Cytokeratin 13 Mouse Monoclonal Antibody (KRT13/2213).




Fig. 2: Formalin-fixed, paraffin-embedded human Prostate Carcinoma stained with Cytokeratin 13 Mouse Monoclonal Antibody (KRT13/2213).




Fig. 3: Flow Cytometric Analysis of HeLa cells using Cytokeratin 13 Mouse Monoclonal Antibody (KRT13/2213) followed by Goat anti-Mouse IgG-CF488 (Blue); Isotype Control (Red).




Fig. 4: SDS-PAGE Analysis of Purified Cytokeratin 13 Mouse Monoclonal Antibody (KRT13/2213). Confirmation of Purity and Integrity of Antibody.