

## 36-2737: Anti-SMAD4 (Pancreatic Adenocarcinoma Marker) Monoclonal Antibody(Clone: SMAD4/2524)

<b>Clonality :</b>	Monoclonal
<b>Clone Name :</b>	SMAD4/2524
<b>Application :</b>	ELISA
<b>Reactivity :</b>	Human
<b>Gene :</b>	STAT6
<b>Gene ID :</b>	4089
<b>Uniprot ID :</b>	Q13485
<b>Alternative Name :</b>	Deleted in Pancreatic Carcinoma 4 (DPC4); JIP; MAD homolog 4 (MADH4); Mothers against decapentaplegic homolog 4 (MADH4); Mothers against DPP homolog 4; MYHRS; SMAD4
<b>Isotype :</b>	Mouse IgG2a, kappa
<b>Immunogen Information :</b>	Recombinant full-length human STAT6 protein

### Description

Signaling from the ligand-activated membrane receptor serine/threonine kinases to nuclear targets is mediated by a set of evolutionarily conserved proteins known as DPC4. Upon ligand binding, the receptors of the TGF- family phosphorylate SMAD proteins (SMAD1 and SMAD2). These proteins then move into the nucleus, where they activate transcription. To carry out this function, the receptor activated SMAD1 and 2 require association with the product of deleted in pancreatic carcinoma, locus 4 (DPC4), also known as SMAD4. SMAD4/DPC4 is also implicated as a tumor suppressor, since it is inactivated in more than half of pancreatic carcinomas and to a lesser extent in a variety of other cancers. SMAD4 is absent in approximately 80% of pancreatic adenocarcinoma, but rarely in endometrial, colorectal, ovarian, lung, breast adenocarcinomas, and malignant melanom. SMAD4 is an important marker for confirming a diagnosis of pancreatic adenocarcinoma. Patients with pancreatic adenocarcinomas with SMAD4 protein expression had significantly longer survival than SMAD4 negative patients.

### Product Info

<b>Amount :</b>	20 µg / 100 µg
<b>Content :</b>	200 µg/ml of Ab Purified from Bioreactor Concentrate by Protein A/G. Prepared in 10mM PBS with 0.05% BSA & 0.05% azide. Also available WITHOUT BSA & azide at 1.0mg/ml.
<b>Storage condition :</b>	Antibody with azide - store at 2 to 8°C. Antibody without azide - store at -20 to -80°C. Antibody is stable for 24 months. Non-hazardous.

### Application Note

ELISA (For coating use Ab at 1-5ug/ml, order Ab without BSA);,




Fig. 1: SDS-PAGE Analysis Purified SMAD4 Mouse Monoclonal Antibody (SMAD4/2524). Confirmation of Purity and Integrity of Antibody.