

36-2764: Anti-Mammaglobin (SCGB2A2) (Breast Cancer Marker) Monoclonal Antibody(Clone: SPM518)

Clonality :	Monoclonal
Clone Name :	SPM518
Application :	WB,IHC
Reactivity :	Human
Gene :	SCGB2A2
Gene ID :	4250
Uniprot ID :	Q13296
Alternative Name :	Mammaglobin-1 (MGB1); Mammaglobin-A; Secretoglobin family 2A member 2 (SCGB2A2); SG2A2; ÅµgB2
Isotype :	Mouse IgG1, kappa
Immunogen Information :	Recombinant full-length human Mammaglobin (SCGB2A2) protein

Description

Mammaglobin is a 93 amino acid glycoprotein with homology to other secretoglobin-uteroglobin family members. It was originally identified as a breast cancer restricted biomarker by differential screening. Mammaglobin related to secretoglobin family that includes human uteroglobin and lipophilin. Mammaglobin antibody stains cytoplasm of normal breast epithelial cells as well as primary and metastatic breast carcinomas. Mammaglobin expression is absent in prostate, kidney, colon, rectum, small intestine, stomach, pancreas, lung, and thyroid tissues. Mammaglobin may be used as part of an immunohistochemical panel for determination of metastatic breast carcinoma and tumor of unknown primary origin.

Product Info

Amount :	20 µg / 100 µg
Content :	200 µg/ml of Ab Purified from Bioreactor Concentrate by Protein A/G. Prepared in 10mM PBS with 0.05% BSA & 0.05% azide. Also available WITHOUT BSA & azide at 1.0mg/ml.
Storage condition :	Antibody with azide - store at 2 to 8°C. Antibody without azide - store at -20 to -80°C. Antibody is stable for 24 months. Non-hazardous.

Application Note

Western Blot (1-2ug/ml); Immunohistochemistry (Formalin-fixed) (1-2ug/ml for 30 min at RT)(Staining of formalin-fixed tissues requires heating tissue sections in 10mM Tris with 1mM EDTA, pH 9.0, for 45 min at 95°C followed by cooling at RT for 20 minutes);




Fig. 1: Formalin-fixed, paraffin-embedded human Breast Carcinoma stained with Monospecific Mouse Monoclonal Antibody (SPM518) to Mammaglobin.




Fig. 2: SDS-PAGE Analysis of Purified Monospecific Mouse Monoclonal Antibody (SPM518) to Mammaglobin. Confirmation of purity and integrity.




Fig. 3: Western Blot Analysis of Human MCF-7, T47D and MDA-231 cell lysate using Monospecific Mouse Monoclonal Antibody (SPM518) to Mammaglobin.




Fig. 4: Analysis of Protein Array containing >19,000 full-length human proteins using Mammaglobin (SCGB2A2) Mouse Monoclonal Antibody (SPM518) Z- and S- Score: The Z-score represents the strength of a signal that a monoclonal antibody (MAb) (in combination with a fluorescently-tagged anti-IgG secondary antibody) produces when binding to a particular protein on the HuProt™ array. Z-scores are described in units of standard deviations (SD's) above the mean value of all signals generated on that array. If targets on HuProt™ are arranged in descending order of the Z-score, the S-score is the difference (also in units of SD's) between the Z-score. S-score therefore represents the relative target specificity of a MAb to its intended target. A MAb is considered to specific to its intended target, if the MAb has an S-score of at least 2.5. For example, if a MAb binds to protein X with a Z-score of 43 and to protein Y with a Z-score of 14, then the S-score for the binding of that MAb to protein X is equal to 29.