

## 36-2777: Anti-CD10 (Membrane Metalloendopeptidase) Monoclonal Antibody(Clone: FR4D11)

<b>Clonality :</b>	Monoclonal
<b>Clone Name :</b>	FR4D11
<b>Application :</b>	FACS,IF
<b>Reactivity :</b>	Human
<b>Gene :</b>	MME
<b>Gene ID :</b>	4311
<b>Uniprot ID :</b>	P08473
<b>Alternative Name :</b>	Atriopeptidase; Common acute lymphocytic leukemia antigen (CALLA); Enkephalinase (EPN); gp100; Membrane metalloendopeptidase (MME); Neprilysin; Neutral endopeptidase (NEP); Skin fibroblast elastase (SFE)
<b>Isotype :</b>	Mouse IgG1, kappa
<b>Immunogen Information :</b>	Raji cells

### Description

Recognizes a 100kDa glycoprotein, identified as CD10, also known as Common Acute Lymphocytic Leukemia Antigen (CALLA). It is a cell surface enzyme with neutral metalloendopeptidase activity, which inactivates a variety of biologically active peptides. CD10 is expressed on the cells of lymphoblastic, Burkitt's, and follicular germinal center lymphomas, and on cells from patients with chronic myelocytic leukemia (CML). It is also expressed on the surface of normal early lymphoid progenitor cells, immature B cells within adult bone marrow and germinal center B cells within lymphoid tissue. CD10 is also present on breast myoepithelial cells, bile canaliculi, fibroblasts, with especially high expression on the brush border of kidney and gut epithelial cells.

### Product Info

<b>Amount :</b>	20 µg / 100 µg
<b>Content :</b>	200 µg/ml of Ab Purified from Bioreactor Concentrate by Protein A/G. Prepared in 10mM PBS with 0.05% BSA & 0.05% azide. Also available WITHOUT BSA & azide at 1.0mg/ml.
<b>Storage condition :</b>	Antibody with azide - store at 2 to 8°C. Antibody without azide - store at -20 to -80°C. Antibody is stable for 24 months. Non-hazardous.

### Application Note

Flow Cytometry (1-2ug/million cells); Immunofluorescence (1-2ug/ml);




Fig. 1: SDS-PAGE Analysis Purified CD10 Mouse Monoclonal Antibody (FR4D11). Confirmation of Integrity and Purity of Antibody




Fig. 2: Flow Cytometric Analysis of Ramos cells using CD10 Mouse Monoclonal Antibody (FR4D11) followed by goat anti-Mouse IgG-CF488 (Blue); Isotype control (Red).




Fig. 3: Immunofluorescence staining of Ramos cells using CD10 Mouse Monoclonal Antibody (FR4D11) followed by goat anti-Mouse IgG conjugated to CF488 (green). Nuclei are stained with Reddot.