

9853 Pacific Heights Blvd. Suite D. San Diego, CA 92121, USA Tel: 858-263-4982

Email: info@abeomics.com

36-2789: Anti-MMP3 (Marker of Metastasis and Rheumatoid Arthritis) Monoclonal Antibody(Clone: MMP3/1730)

Clonality: Monoclonal Clone Name: MMP3/1730

Application: IHC
Reactivity: Human
Gene: MMP3
Gene ID: 4314
Uniprot ID: P08254

Alternative Name: CHDS6; Matrix metalloproteinase 3; Matrix metalloproteinase-3; MMP3; Proteoglycanase; SL1;

STMY1; STR1; Stromelysin-1; Transin 1

Isotype: Mouse IgG1, lambda

Immunogen Information: Recombinant full-length human MMP3 protein

Description

The matrix metalloproteinases (MMP) are a family of peptidase enzymes responsible for the degradation of extracellular matrix components, including collagen, gelatin, fibronectin, laminin and proteoglycan. Transcription of MMP genes is differentially activated by phorbol ester, lipopolysaccharide (LPS) or staphylococcal enterotoxin B (SEB). MMP catalysis requires both calcium and zinc. MMP-3, MMP-10 and MMP-11 (also designated stromelysin-1, 2 and 3, respectively) activate procollagenase. MMP-3 activation of procollagenase can occur via two pathways. Direct activation by MMP-3 is slow and activation by MMP-3 in conjunction with tissue or plasma proteinases is rapid. MMP-10 is expressed in small intestine, and at lower levels in lung and heart. MMP-11 is specifically expressed in stromal cells of breast carcinomas and contributes to epithelial cell malignancies.

Product Info

Amount: 20 μg / 100 μg

Content: 200 μg/ml of Ab Purified by Protein A. Prepared in 10mM PBS with 0.05% BSA & 0.05% azide.

Also available WITHOUT BSA & azide at 1.0mg/ml.

Storage condition : Antibody with azide - store at 2 to 8°C. Antibody without azide - store at -20 to -80°C. Antibody

is stable for 24 months. Non-hazardous.

Application Note

Immunohistochemistry (Formalin-fixed) (1-2ug/ml for 30 minutes at RT),(Staining of formalin-fixed tissues requires heating tissue sections in 10mM Tris with 1mM EDTA, pH 9.0, for 45 min at 95 °C followed by cooling at RT for 20 minutes),




Fig. 1: SDS-PAGE Analysis of Purified MMP3 Mouse Monoclonal Antibody (MMP3/1730). Confirmation of Integrity and Purity of Antibody.