

9853 Pacific Heights Blvd. Suite D. San Diego, CA 92121, USA Tel: 858-263-4982

Email: info@abeomics.com

36-2794: Anti-MMP9 (Matrix Metalloproteinase 9) Monoclonal Antibody(Clone: 2C3)

Clonality: Monoclonal

Clone Name: 2C3
Application: IHC
Reactivity: Human
Gene: MMP9
Gene ID: 4318
Uniprot ID: P14780

82 kDa matrix metalloproteinase-9; 92 kDa type IV collagenase; CLG 4B; CLG4B; Collagenase

Alternative Name: Type 4 beta; Collagenase type IV 92 KD; EC 3.4.24.35; Gelatinase 92 KD; Gelatinase B;

Gelatinase beta; GELB; Macrophage gelatinase; MANDP2;

Isotype: Mouse IgG1, kappa

Immunogen Information: A synthetic peptide corresponding to amino acids 603-614 of human MMP-9

Description

The matrix metalloproteinases (MMP) are a family of peptidase enzymes responsible for the degradation of extracellular matrix components, including collagen, gelatin, fibronectin, laminin and proteoglycan. Transcription of MMP genes is differentially activated by phorbol ester, lipopolysaccharide (LPS) or staphylococcal enterotoxin B (SEB). MMP catalysis requires both calcium and zinc. MMP-9 (also designated 92 kDa type IV collagenase or gelatinase B) has been shown to degrade bone collagens in concert with MMP-1 (also designated interstitial collagenase, fibroblast collagenase or collagenase-1), and cysteine proteases and may play a role in bone osteoclastic resorption. MMP-1 is down-regulated by p53, and abnormality of p53 expression may contribute to joint degradation in rheumatoid arthritis by regulating MMP-1 expression.

Product Info

Amount : 20 μg / 100 μg

Content: 200 μg/ml of Ab Purified by Protein A. Prepared in 10mM PBS with 0.05% BSA & 0.05% azide.

Also available WITHOUT BSA & azide at 1.0mg/ml.

Storage condition : Antibody with azide - store at 2 to 8°C. Antibody without azide - store at -20 to -80°C. Antibody

is stable for 24 months. Non-hazardous.

Application Note

Immunohistochemistry (Formalin-fixed) (1-2ug/ml for 30 minutes at RT)(Staining of formalin-fixed tissues requires heating tissue sections in 10mM Tris with 1mM EDTA, pH 9.0, for 45 min at 95&deaC followed by cooling at RT for 20 minutes):




Fig. 1: Formalin-fixed, paraffin-embedded human Spleen stained with MMP9 Mouse Monoclonal Antibody (2C3).



9853 Pacific Heights Blvd. Suite D. San Diego, CA 92121, USA Tel: 858-263-4982

Email: info@abeomics.com




Fig. 2: SDS-PAGE Analysis of Purified MMP9 Mouse Monoclonal Antibody (2C3). Confirmation of Integrity and Purity of Antibody