

## 36-2840: Anti-CD56 / NCAM1 / NKH1 (Neuronal Cell Marker) Monoclonal Antibody(Clone: ERIC-1)

<b>Clonality :</b>	Monoclonal
<b>Clone Name :</b>	ERIC-1
<b>Application :</b>	IHC
<b>Reactivity :</b>	Human
<b>Gene :</b>	NCAM1
<b>Gene ID :</b>	4684
<b>Alternative Name :</b>	NCAM, Leu-19, NKH1, MSK39, NCAM120, NCAM140, NCAM180, Neural Cell Adhesion Molecule
<b>Isotype :</b>	Mouse IgG1, kappa
<b>Immunogen Information :</b>	Recombinant human NCAM1 protein

### Description

This MAb reacts with an extracellular domain (close to transmembrane) of CD56/NCAM. Three isoforms of neural cell adhesion molecule (NCAM) are produced by differential splicing of the RNA transcript from a single gene. The 135kDa isoform is the basic molecule, which is glycosylated or sialylated to produce the mature species. Anti-CD56 recognizes two proteins of the neural cell adhesion molecule, the basic molecule expressed on most neuroectodermally derived tissues and neoplasms (e.g. retinoblastoma, medulloblastomas, astrocytomas, neuroblastomas, and small cell carcinomas). It is also expressed on some mesodermally derived tumors (rhabdomyosarcoma). Anti-CD56 plays an important role in the diagnosis of nodal and nasal NK/T-cell lymphomas.

### Product Info

<b>Amount :</b>	20 µg / 100 µg
<b>Content :</b>	200 µg/ml of Ab Purified from Bioreactor Concentrate by Protein A/G. Prepared in 10mM PBS with 0.05% BSA & 0.05% azide. Also available WITHOUT BSA & azide at 1.0mg/ml.
<b>Storage condition :</b>	Antibody with azide - store at 2 to 8°C. Antibody without azide - store at -20 to -80°C. Antibody is stable for 24 months. Non-hazardous.

### Application Note

Immunohistochemistry (Formalin-fixed) (0.1-0.2µg/ml for 30 min at RT)(Staining of formalin-fixed tissues requires heating tissue sections in 10mM Tris with 1mM EDTA, pH 9.0, for 45 min at 95°C followed by cooling at RT for 20 minutes);




Fig. 1: Formalin-fixed, paraffin-embedded human Colon stained with CD56 Mouse Monoclonal Antibody (ERIC-1).




Fig. 2: Formalin-fixed, paraffin-embedded human Colon stained with CD56 Mouse Monoclonal Antibody (ERIC-1).




Fig. 3: SDS-PAGE Analysis Purified CD56 Mouse Monoclonal Antibody (ERIC-1). Confirmation of Purity and Integrity of Antibody.