

## 36-2889: Anti-Langerin / CD207 (Marker of Langerhans Cells) Monoclonal Antibody(Clone: rLGRN/1821)

<b>Clonality :</b>	Monoclonal
<b>Clone Name :</b>	rLGRN/1821
<b>Application :</b>	ELISA
<b>Reactivity :</b>	Human
<b>Gene :</b>	CD207
<b>Gene ID :</b>	50489
<b>Uniprot ID :</b>	Q9UJ71
<b>Alternative Name :</b>	C-type lectin domain family 4 member K; CD207; CLEC4K; Langerhans cell specific c type lectin; Langerin
<b>Isotype :</b>	Mouse IgG1, kappa
<b>Immunogen Information :</b>	Recombinant fragment (around aa74-213) of human Langerin protein (exact sequence is proprietary)

### Description

Langerhans cells (LCs) are a subset of immature dendritic cells (DCs) that specifically localize in the epidermis and other mucosal epithelia. Epidermal LCs possess strong immuno-stimulatory capacity and play a central role in the initiation and regulation of immune responses. Langerin (CD207) is a Ca<sup>2+</sup>-dependent, C-type lectin domain containing, type II transmembrane protein that induces epidermal LCs to differentiate into Birbeck granules (BG). BGs are organelles with superimposing and zipper membranes that influence proper class I type antigen presentation to the circulating T cells. Human spleen, lymph node, thymus, liver, lung and heart express Langerin protein.

### Product Info

<b>Amount :</b>	20 µg / 100 µg
<b>Content :</b>	200 µg/ml of Ab Purified from Bioreactor Concentrate by Protein A/G. Prepared in 10mM PBS with 0.05% BSA & 0.05% azide. Also available WITHOUT BSA & azide at 1.0mg/ml.
<b>Storage condition :</b>	Antibody with azide - store at 2 to 8°C. Antibody without azide - store at -20 to -80°C. Antibody is stable for 24 months. Non-hazardous.

### Application Note

ELISA (Use Ab at 2-4ug/ml for coating) (Order Ab without BSA);




Fig. 1: SDS-PAGE Analysis Purified Langerin Recombinant Mouse Monoclonal Antibody (rLGRN/1821). Confirmation of Integrity and Purity of Antibody.