

## 36-2901: Anti-FOXP3 (Forkhead Box Protein P3) / Scurfin Monoclonal Antibody(Clone: 3G3)

<b>Clonality :</b>	Monoclonal
<b>Clone Name :</b>	3G3
<b>Application :</b>	FACS,IF
<b>Reactivity :</b>	Human, Mouse
<b>Gene :</b>	FOXP3
<b>Gene ID :</b>	50943
<b>Uniprot ID :</b>	Q9ZS1
<b>Alternative Name :</b>	JM2, AIID, DIETER, IPEX, MGC141961, MGC141963, PIDX, XPID, FOXP3delta7, forkhead box protein P3, immune dysregulation, polyendocrinopathy, enteropathy, X-linked, immunodeficiency, polyendocrinopathy, enteropathy, X-linked, scurfin
<b>Isotype :</b>	Mouse IgG1, kappa
<b>Immunogen Information :</b>	Full-length His-tagged Mouse FOXP3 protein

### Description

Recognizes a protein of 47-55kDa, which is identified as FOXP3. Its precise epitope is not known, but it has been mapped to the N-terminal portion (amino acids 2-136) of the protein. The FOX family of transcription factors is a large group of proteins that share a common DNA binding domain termed a winged-helix or forkhead domain. During early development, FOXP1 and FOXP2 are expressed abundantly in the lung, with lower levels of expression in neural, intestinal and cardiovascular tissues, where they act as transcription repressors. FOXP1 is widely expressed in adult tissues, while neoplastic cells often exhibit a dramatic change in expression level or localization of FOXP1. Mutations in FOXP3 gene cause IPEX, a fatal, X-linked inherited disorder characterized by immune dysregulation. The FOXP3 protein is essential for normal immune homeostasis. Specifically, FOXP3 represses transcription through a DNA binding forkhead domain, thereby regulating T cell activation.

### Product Info

<b>Amount :</b>	20 µg / 100 µg
<b>Content :</b>	200 µg/ml of Ab Purified from Bioreactor Concentrate by Protein A/G. Prepared in 10mM PBS with 0.05% BSA & 0.05% azide. Also available WITHOUT BSA & azide at 1.0mg/ml.
<b>Storage condition :</b>	Antibody with azide - store at 2 to 8°C. Antibody without azide - store at -20 to -80°C. Antibody is stable for 24 months. Non-hazardous.

### Application Note

Flow Cytometry (1-2ug/million cells); Immunofluorescence (1-2ug/ml);




Fig. 1: SDS-PAGE Analysis Purified FOXP3 Mouse Monoclonal Antibody (3G3). Confirmation of Purity and Integrity of Antibody.