

9853 Pacific Heights Blvd. Suite D. San Diego, CA 92121, USA Tel: 858-263-4982

Email: info@abeomics.com

## 36-2932: Anti-Prohibitin (Mitochondrial Marker) Monoclonal Antibody(Clone: PHB/1882)

Clonality: Monoclonal
Clone Name: PHB/1882
Application: WB,IHC
Reactivity: Human
Gene: PHB
Gene ID: 5245
Uniprot ID: P35232

Alternative Name: Epididymis luminal protein 215; Epididymis secretory sperm binding protein Li 54e; HEL 215;

HEL S 54e; PHB; PHB1; Prohibitin

**Isotype:** Mouse IgG1, kappa

Immunogen Information: Recombinant human PHB protein fragment (aa167-261) (exact sequence is proprietary)

## **Description**

Recognizes a protein of 30kDa which is identified as Prohibitin, an evolutionarily conserved protein with homologues found in yeast to man. It is located in the inner membrane of mitochondria. Althoµgh prohibitin mRNA and protein expression occurs throµghout the cell cycle, maximum levels are detected during the G1/S phase transition and minimum levels are seen in S phase and the G2/mitosis boundary. Prohibitin is located exclusively in the mitochondria with the highest concentration on the inner membrane. Prohibitin is an ideal mitochondrial marker. It shows antiproliferative activity and has been proposed to play a role in normal cell cycle regulation, replicative senescence, cellular immortalization, and tumor suppression.

## **Product Info**

Amount :  $20 \mu g / 100 \mu g$ 

Content: 200 µg/ml of Ab Purified from Bioreactor Concentrate by Protein A/G. Prepared in 10mM PBS

with 0.05% BSA & 0.05% azide. Also available WITHOUT BSA & azide at 1.0mg/ml.

**Storage condition :** Antibody with azide - store at 2 to 8°C. Antibody without azide - store at -20 to -80°C. Antibody

is stable for 24 months. Non-hazardous.

## **Application Note**

Western Blot (1-2ug/ml);,Immunohistochemistry (Formalin-fixed) (1-2ug/ml for 30 minutes at RT),(Staining of formalin-fixed tissues requires heating tissue sections in 10mM Tris with 1mM EDTA, pH 9.0, for 45 min at 95&degC followed by cooling at RT for 20 minutes);




Fig. 1: Formalin-fixed, paraffin-embedded human Liver stained with Prohibitin Mouse Monoclonal Antibody (PHB/1882).



9853 Pacific Heights Blvd. Suite D. San Diego, CA 92121, USA Tel: 858-263-4982

Email: info@abeomics.com




Fig. 2: Western Blot Analysis of (1) A549 (2) K562 (3) A431 & (4) MCF-7 cell lysates using Prohibitin Mouse Monoclonal Antibody (PHB/1882).




Fig. 3: Western Blot Analysis of human Ramos, HEK293, K562, Jurkat & HepG2 cell lysates using Prohibitin Mouse Monoclonal Antibody (PHB/1882).




Fig. 4: SDS-PAGE Analysis Purified Prohibitin Mouse Monoclonal Antibody (PHB/1882). Confirmation of Purity and Integrity of Antibody.