

9853 Pacific Heights Blvd. Suite D. San Diego, CA 92121, USA Tel: 858-263-4982

Email: info@abeomics.com

## 36-2940: Anti-Alpha-1-Antitrypsin (SERPINA1) (Hepatocellular & Histiocytic Marker) Monoclonal Antibody(Clone: AAT/1378)

Clonality: Monoclonal
Clone Name: AAT/1378
Application: WB,IHC
Reactivity: Human
Gene: SERPINA1
Gene ID: 5265
Uniprot ID: P01009

A1AT; AAT; Alpha 1-antitrypsin; Alpha-1 protease inhibitor; Alpha-1-antiproteinase; alpha1

Alternative Name: proteinase inhibitor; Alpha1AT; Dom1; Serine (or cysteine) proteinase inhibitor clade A

member 1; Serine protease inhibitor 1-1; Serine protease inhibitor A1a; Serpin A1a; Serpin peptidase inhibitor clade A member 1; Serpina1; Short peptide from AAT; SPAAT; Spi1-1

**Isotype:** Mouse IgG1, kappa

Immunogen Information: Recombinant human Alpha-1-Antitrypsin (AAT) protein fragment

## **Description**

It recognizes a protein of 54kDa, which is identified antitrypsin (AAT). The immunohistochemical staining of AAT is useful in identification of benign and malignant hepatic tumors and yolk sac carcinomas. Positive staining for AAT is also used in detection of benign and malignant lesions of histiocytic nature. This antibody is may also useful tool in the screening of patients with cryptogenic cirrhosis or other forms of liver disease with fibrosis of uncertain origin.

## **Product Info**

**Amount:** 20 μg / 100 μg

Content: 200 µg/ml of Ab Purified from Bioreactor Concentrate by Protein A/G. Prepared in 10mM PBS

with 0.05% BSA & 0.05% azide. Also available WITHOUT BSA & azide at 1.0mg/ml.

. ... Antibody with azide - store at 2 to 8°C. Antibody without azide - store at -20 to -80°C. Antibody

Storage condition: Altibody with azide - store at 2 to 8 C. 7 is stable for 24 months. Non-hazardous.

## **Application Note**

Western Blot (1-2ug/ml); Immunohistochemistry (Formalin-fixed) (1-2ug/ml for 30 minutes at RT)(Staining of formalin-fixed tissues requires heating tissue sections in 10mM Tris with 1mM EDTA, pH 9.0, for 45 min at 95&degC followed by cooling at RT for 20 minutes):




Fig. 1: Formalin-fixed, paraffin-embedded human Tonsil stained with Alpha-1-Antitrypsin Mouse Monoclonal Antibody (AAT/1378).



9853 Pacific Heights Blvd. Suite D. San Diego, CA 92121, USA Tel: 858-263-4982 Email: info@abeomics.com




Fig. 2: Western Blot Analysis of 1) recombinant Alpha-1-Antitrypsin 2) Jurkat & 3) A549 cell lysate using Alpha-1-Antitrypsin Mouse Monoclonal Antibody (AAT/1378).




Fig. 3: SDS-PAGE Analysis Purified Alpha-1-Antitrypsin Mouse Monoclonal Antibody (AAT/1378). Confirmation of Integrity and Purity of Antibody