

36-2943: Anti-PTEN (Tumor Suppressor Oncoprotein) Monoclonal Antibody(Clone: PTEN/2110)

Clonality :	Monoclonal
Clone Name :	PTEN/2110
Application :	IHC
Reactivity :	Human
Gene :	PTEN
Gene ID :	5278
Uniprot ID :	P60484
Alternative Name :	BZS; DEC; GLM2; MHAM; Mutated in Multiple Advanced Cancers 1 (MMAC1); MMAC1 phosphatase and tensin homolog deleted on chromosome 10; Phosphatase and Tensin Homolog; Phosphatase and tensin like protein; Phosphatidylinositol 3,4,5-trisphosphate 3-phosphatase and dual-specificity protein phosphatase; PTEN; PTEN1; TEP1
Isotype :	Mouse IgG2b, kappa
Immunogen Information :	Recombinant full-length human PTEN protein

Description

PTEN (phosphatidylinositol-3,4,5-trisphosphate 3-phosphatase) contains a tensin like domain as well as a catalytic domain similar to that of the dual specificity protein tyrosine phosphatases. It was identified as a tumor suppressor that is mutated in a large number of cancers, including sporadic brain, breast, kidney, and prostate cancers.

Product Info

Amount :	20 µg / 100 µg
Content :	200 µg/ml of Ab Purified from Bioreactor Concentrate by Protein A/G. Prepared in 10mM PBS with 0.05% BSA & 0.05% azide. Also available WITHOUT BSA & azide at 1.0mg/ml.
Storage condition :	Antibody with azide - store at 2 to 8°C. Antibody without azide - store at -20 to -80°C. Antibody is stable for 24 months. Non-hazardous.

Application Note

Immunohistochemistry (Formalin-fixed) (1-2ug/ml for 30 minutes at RT)(Staining of formalin-fixed tissues requires heating tissue sections in 10mM Tris with 1mM EDTA, pH 9.0, for 45 min at 95°C followed by cooling at RT for 20 minutes);




Fig. 1: Formalin-fixed, paraffin-embedded human Prostate Carcinoma stained with PTEN Mouse Monoclonal Antibody (PTEN/2110).




Fig. 2: Formalin-fixed, paraffin-embedded human Renal Cell Carcinoma stained with PTEN Mouse Monoclonal Antibody (PTEN/2110).




Fig. 3: SDS-PAGE Analysis Purified PTEN Mouse Monoclonal Antibody (PTEN/2110). Confirmation of Purity and Integrity of Antibody.




Fig. 4: Analysis of Protein Array containing more than 19,000 full-length human proteins using PTEN Mouse Monoclonal Antibody (PTEN/2110). Z- and S- Score: The Z-score represents the strength of a signal that a monoclonal antibody (MAb) (in combination with a fluorescently-tagged anti-IgG secondary antibody) produces when binding to a particular protein on the HuProt™ array. Z-scores are described in units of standard deviations (SD's) above the mean value of all signals generated on that array. If targets on HuProt™ are arranged in descending order of the Z-score, the S-score is the difference (also in units of SD's) between the Z-score. S-score therefore represents the relative target specificity of a MAb to its intended target. A MAb is considered to specific to its intended target, if the MAb has an S-score of at least 2.5. For example, if a MAb binds to protein X with a Z-score of 43 and to protein Y with a Z-score of 14, then the S-score for the binding of that MAb to protein X is equal to 29.