

## 36-2955: Anti-ACTH (Adrenocorticotrophic Hormone) (C-Terminus) Monoclonal Antibody(Clone: rCLIP/1418)

<b>Clonality :</b>	Monoclonal
<b>Clone Name :</b>	rCLIP/1418
<b>Application :</b>	ELISA
<b>Reactivity :</b>	Human
<b>Gene :</b>	POMC
<b>Gene ID :</b>	5443
<b>Uniprot ID :</b>	P01189
<b>Alternative Name :</b>	Adrenocorticotropin; alpha or beta or gamma Melanocyte Stimulating Hormone (MSH) or Melanotropin; beta-Endorphin; beta or gamma Lipotropin (LPH); CLIP; Met Enkephalin; POC; POMC
<b>Isotype :</b>	Mouse IgG1, kappa
<b>Immunogen Information :</b>	Synthetic peptide corresponding to aa25-39 of human ACTH (NGAEDESAEAFPLEF)

### Description

ACTH (same as Corticotropin) is a 39 amino acid active peptide produced by the anterior pituitary. This MAb is specific to CLIP (aa25-39 of ACTH); does not react with Synacthen (aa1-24 of ACTH). POMC (pro-opiomelanocortin or corticotropin-lipotropin) is a 267 amino acid polypeptide hormone precursor that goes through extensive, tissue-specific posttranslational processing by convertases. POMC is cleaved into ten hormone chains named NPP, ACTH, alpha-MSH (Melanocyte Stimulating Hormone), beta-MSH, gamma-MSH, CLIP (corticotropin-like intermediary peptide), Lipotropin-beta, Lipotropin-gamma, beta-endorphin and Met-enkephalin. ACTH is also produced by cells of immune system (T-cells, B-cells, and macrophages) in response to stimuli associated with stress. Anti-ACTH is a useful marker in classification of pituitary tumors and the study of pituitary disease. It reacts with ACTH-producing cells (corticotrophs). It also may react with other tumors (e.g. some small cell carcinomas of the lung) causing paraneoplastic syndromes by secreting ACTH. '

### Product Info

<b>Amount :</b>	20 µg / 100 µg
<b>Content :</b>	200 µg/ml of Ab Purified from Bioreactor Concentrate by Protein A/G. Prepared in 10mM PBS with 0.05% BSA & 0.05% azide. Also available WITHOUT BSA & azide at 1.0mg/ml.
<b>Storage condition :</b>	Antibody with azide - store at 2 to 8°C. Antibody without azide - store at -20 to -80°C. Antibody is stable for 24 months. Non-hazardous.

### Application Note

ELISA (For coating, order Ab without BSA);




Fig. 1: SDS-PAGE Analysis Purified ACTH Recombinant Mouse Monoclonal Antibody (rCLIP/1418). Confirmation of Purity and Integrity of Antibody.