

36-2976: Anti-OCT-2 (POU2F2) (B-Cell Marker) Monoclonal Antibody(Clone: 86839)

Clonality :	Monoclonal
Clone Name :	Oct2-2137
Application :	ELISA,WB,IHC
Reactivity :	Human
Gene :	POU2F2
Gene ID :	5452
Uniprot ID :	P09086
Alternative Name :	Lymphoid-restricted immunoglobulin octamer-binding protein NF-A2; Oct-2; Octamer-binding protein 2; Octamer-binding transcription factor 2; OTF-2; POU domain class 2 transcription factor 2; POU2F2
Isotype :	Mouse IgG1, kappa
Immunogen Information :	Recombinant fragment of human OCT2 protein (around aa 112-297) (exact sequence is proprietary)

Description

Oct-2 is a transcription factor of the POU homeo-domain family that binds to the Ig gene octamer sites, regulating B-cell-specific genes. Oct-2 expression can be used as a marker of B-cell lineage and differentiation. Germinal center B-cells, mantle B-cells, monocytoid B-cells, and plasma cells show high level expression of Oct-2. Additionally, mantle cell lymphoma, follicular lymphoma, marginal zone lymphoma, plasmacytoma, Burkitt lymphoma, diffuse large cell lymphoma, diffuse large B-cell lymphoma, Hodgkin lymphoma display increased expression of Oct-2. Several studies of Oct-2 expression have shown a low level expression in pre-B, T-cell, myelomonocytic, and epithelial cell lines, whereas all mature B-cell lines have high levels of expression. In spite of scanty evidence for Oct-2 expression in T cells, it is believed that this factor participates in transcriptional regulation during T-cell activation.

Product Info

Amount :	20 µg / 100 µg
Content :	200 µg/ml of Ab Purified from Bioreactor Concentrate by Protein A/G. Prepared in 10mM PBS with 0.05% BSA & 0.05% azide. Also available WITHOUT BSA & azide at 1.0mg/ml.
Storage condition :	Antibody with azide - store at 2 to 8°C. Antibody without azide - store at -20 to -80°C. Antibody is stable for 24 months. Non-hazardous.

Application Note

ELISA (For coating, order antibody without BSA); Western Blot (1-2ug/ml); ,Immunohistochemistry (Formalin-fixed) (1-2ug/ml for 30 minutes at RT),(Staining of formalin-fixed tissues requires heating tissue sections in 10mM Tris with 1mM EDTA, pH 9.0, for 45 min at 95 °C followed by cooling at RT for 20 minutes),




Fig. 1: Formalin-fixed, paraffin-embedded human Lymph Node stained with Oct-2 Mouse Monoclonal Antibody (OCT2/2137).




Fig. 2: Formalin-fixed, paraffin-embedded human Tonsil stained with Oct-2 Mouse Monoclonal Antibody (OCT2/2137).




Fig. 3: Western Blot Analysis Ramos cell lysate using Oct-2 Mouse Monoclonal Antibody (OCT2/2137).




Fig. 4: Western Blot Analysis (A) Recombinant Protein (B) Daudi cell lysate Oct-2 Mouse Monoclonal Antibody (OCT2/2137).




Fig. 5: SDS-PAGE Analysis Purified Oct-2 Mouse Monoclonal Antibody (OCT2/2137). Confirmation of Integrity and Purity of Antibody.




Fig. 6: Analysis of Protein Array containing more than 19,000 full-length human proteins using Oct-2 Mouse Monoclonal Antibody (OCT2/2137) Z- and S- Score: The Z-score represents the strength of a signal that a monoclonal antibody (MAB) (in combination with a fluorescently-tagged anti-IgG secondary antibody) produces when binding to a particular protein on the HuProt™ array. Z-scores are described in units of standard deviations (SD's) above the mean value of all signals generated on that array. If targets on HuProt™ are arranged in descending order of the Z-score, the S-score is the difference (also in units of SD's) between the Z-score. S-score therefore represents the relative target specificity of a MAB to its intended target. A MAB is considered to specific to its intended target, if the MAB has an S-score of at least 2.5. For example, if a MAB binds to protein X with a Z-score of 43 and to protein Y with a Z-score of 14, then the S-score for the binding of that MAB to protein X is equal to 29.