

9853 Pacific Heights Blvd. Suite D. San Diego, CA 92121, USA Tel: 858-263-4982

Email: info@abeomics.com

## 36-3056: Anti-Parathyroid Hormone (PTH) (N-Terminal) Monoclonal Antibody(Clone: PTH/1717R)

Clonality: Monoclonal Clone Name: PTH/1717R

Application: IHC
Reactivity: Human
Gene: PTH
Gene ID: 5741
Uniprot ID: P01270

Alternative Name: hPTH; Parathormone; Parathyrin; Parathyroid hormone 1 (PTH1); Parathyroid hormone (PTH)

**Isotype:** Rabbit IgG

Immunogen Information: A synthetic peptide around aa 1-34 of human mature-PTH-polypeptide (exact sequence is

proprietary)

## **Description**

Epitope of this MAb maps in the N-terminus of PTH, a hormone produced by the parathyroid gland that regulates the concentration of calcium and phosphorus in extracellular fluid. This hormone elevates blood Ca2+ levels by dissolving the salts in bone and preventing their renal excretion. It is produced in the parathyroid gland as an 84 amino acid single chain polypeptide. It can also be secreted as N-terminal truncated fragments or C-terminal fragments after intracellular degradation, as in case of hypercalcemia. Defects in this gene are a cause of familial isolated hypoparathyroidism (FIH); also called autosomal dominant hypoparathyroidism or autosomal dominant hypocalcemia. FIH is characterized by hypocalcemia and hyperphosphatemia due to inadequate secretion of parathyroid hormone. Symptoms are seizures, tetany and cramps. FIH exist both as autosomal dominant and recessive forms of hypoparathyroidism.

## **Product Info**

**Amount:** 20 μg / 100 μg

**Content:** 200 μg/ml of Ab Purified by Protein A. Prepared in 10mM PBS with 0.05% BSA & 0.05% azide.


Also available WITHOUT BSA & azide at 1.0mg/ml.

**Storage condition :** Antibody with azide - store at 2 to 8°C. Antibody without azide - store at -20 to -80°C. Antibody

is stable for 24 months. Non-hazardous.

## **Application Note**

Immunohistochemistry (Formalin-fixed) (1-2ug/ml for 30 minutes at RT)(Staining of formalin-fixed tissues requires heating tissue sections in 10mM Tris with 1mM EDTA, pH 9.0, for 45 min at 95&degC followed by cooling at RT for 20 minutes);



 $Fig. \ 1: Formalin-fixed, paraffin-embedded \ human \ Parathyroid \ stained \ with \ PTH \ Rabbit \ Recombinant \ Monoclonal \ Antibody \ (PTH/1717R).$ 



9853 Pacific Heights Blvd. Suite D. San Diego, CA 92121, USA Tel: 858-263-4982

Email: info@abeomics.com




Fig. 2: SDS-PAGE Analysis Purified PTH Rabbit Recombinant Monoclonal Antibody (PTH/1717R). Confirmation of Purity and Integrity of Antibody.