

36-3074: Anti-CD45RB (B-Cell Marker) Monoclonal Antibody(Clone: PTPRC/1783R)

Clonality :	Monoclonal
Clone Name :	PTPRC/1783R
Application :	FACS,IF,WB,IHC
Reactivity :	Human
Gene :	PTPRC
Gene ID :	5788
Uniprot ID :	P08575
Alternative Name :	B220, CD45R, GP180, Leukocyte common antigen (LCA), Loc, Ly-5, Lyt-4, Protein tyrosine phosphatase receptor type C (PTPRC), Receptor-type tyrosine-protein phosphatase C, T200 glycoprotein
Isotype :	Rabbit IgG
Immunogen Information :	Recombinant human full-length CD45RB protein

Description

CD45R, also designated CD45 and PTPRC, has been identified as a transmembrane glycoprotein, broadly expressed among hematopoietic cells. Multiple isoforms of CD45R are distributed throughout the immune system according to cell type. These isoforms arise because of alternative splicing of exons 4, 5, and 6. The corresponding protein domains are characterized by the binding of monoclonal antibodies specific for CD45RA (exon 4), CD45RB (exon 5), CD45RC (exon 6) and CD45RO (exons 4 to 6 spliced out). The variation in these isoforms is localized to the extracellular domain of CD45R, while the intracellular domain is conserved. CD45RB is expressed on mature B-lymphocytes and the majority of lymphomas and leukemias of B-cell origin.

Product Info

Amount :	20 µg / 100 µg
Content :	200 µg/ml of Ab Purified by Protein A. Prepared in 10mM PBS with 0.05% BSA & 0.05% azide. Also available WITHOUT BSA & azide at 1.0mg/ml.
Storage condition :	Antibody with azide - store at 2 to 8°C. Antibody without azide - store at -20 to -80°C. Antibody is stable for 24 months. Non-hazardous.

Application Note

Flow Cytometry (1-2ug/million cells); Immunofluorescence (1-2ug/ml); Western Blot (1-2ug/ml); Immunohistochemistry (Formalin-fixed) (1-2ug/ml for 30 minutes at RT),(Staining of formalin-fixed tissues requires heating tissue sections in 10mM Tris with 1mM EDTA, pH 9.0, for 45 min at 95 °C followed by cooling at RT for 20 minutes),




Fig. 1: Formalin-fixed, paraffin-embedded human Tonsil stained with CD45RB Rabbit Recombinant Monoclonal Antibody (PTPRC/1783R).




Fig. 2: Paraformaldehyde-fixed Raji cells stained with CD45RB Rabbit Recombinant Monoclonal Antibody (PTPRC/1783R) followed by Goat anti-rabbit IgG-CF488 (Green). Counterstain is Phalloidin (red).




Fig. 3: Flow Cytometric Analysis of paraformaldehyde-fixed Raji cells using CD45RB Rabbit Recombinant Monoclonal Antibody (PTPRC/1783R) followed by Goat anti-rabbit IgG-CF488 (Blue); Isotype Control (Red).




Fig. 4: Immunofluorescence Analysis of PFA-fixed K562 cells stained with Nucleolin MAb (SPM614) followed by Goat anti-Mouse IgG-CF488 (Green). Membrane is stained with Phalloidin-CF640.




Fig. 5: SDS-PAGE Analysis of Purified CD45RB Rabbit Recombinant Monoclonal Antibody (PTPRC/1783R). Confirmation of Purity and Integrity of Antibody.