

36-3089: Anti-Cyclin D1 (G1-Cyclin & Mantle Cell Lymphoma Marker) Monoclonal Antibody(Clone: CCND1/3548)

Clonality :	Monoclonal
Clone Name :	CCND1/3548
Application :	ELISA
Reactivity :	Human
Gene :	CCND1
Gene ID :	595
Uniprot ID :	P24385
Alternative Name :	B cell CLL/lymphoma 1; B cell leukemia 1; B-cell lymphoma 1 protein; BCL-1 oncogene; CCND1 protein; CCND1/FSTL3 fusion gene; CCND1/IGHG1 fusion gene CCND1/IGLC1 fusion gene; CCND1/PTH fusion gene; G1/S-specific cyclin-D1, Parathyroid adenomatosis 1, PRAD1 oncogene
Isotype :	Mouse IgG2b, kappa
Immunogen Information :	Recombinant fragment of human CCND1 (around aa 115-270) protein (exact sequence is proprietary)

Description

Recognizes a protein of 36kDa, identified as cyclin D1. Cyclin D1, one of the key cell cycle regulators, is a putative proto-oncogene overexpressed in a wide variety of human neoplasms. This antibody neutralizes the activity of cyclin D1 in vivo. About 60% of mantle cell lymphomas (MCL) contain a t(11; 14)(q13; q32) translocation resulting in over-expression of cyclin D1. This antibody is useful in identifying mantle cell lymphomas (cyclin D1 positive) from CLL/SLL and follicular lymphomas (cyclin D1 negative). Occasionally, hairy cell leukemia and plasma cell myeloma weakly express Cyclin D1.

Product Info

Amount :	20 µg / 100 µg
Content :	200 µg/ml of Ab Purified from Bioreactor Concentrate by Protein A/G. Prepared in 10mM PBS with 0.05% BSA & 0.05% azide. Also available WITHOUT BSA & azide at 1.0mg/ml.
Storage condition :	Antibody with azide - store at 2 to 8°C. Antibody without azide - store at -20 to -80°C. Antibody is stable for 24 months. Non-hazardous.

Application Note

ELISA (For coating, order Ab without BSA);




Fig. 1: SDS-PAGE Analysis Purified Cyclin D1 Mouse Monoclonal Antibody (CCND1/3548). Confirmation of Purity and Integrity of Antibody.




Fig. 2: Analysis of Protein Array containing more than 19,000 full-length human proteins using Cyclin D1 Mouse Monoclonal Antibody (CCND1/2593). Z- and S- Score: The Z-score represents the strength of a signal that a monoclonal antibody (Monoclonal Antibody) (in combination with a fluorescently-tagged anti-IgG secondary antibody) produces when binding to a particular protein on the HuProt™ array. Z-scores are described in units of standard deviations (SD's) above the mean value of all signals generated on that array. If targets on HuProt™ are arranged in descending order of the Z-score, the S-score is the difference (also in units of SD's) between the Z-score. S-score therefore represents the relative target specificity of a Monoclonal Antibody to its intended target. A Monoclonal Antibody is considered to specific to its intended target, if the Monoclonal Antibody has an S-score of at least 2.5. For example, if a Monoclonal Antibody binds to protein X with a Z-score of 43 and to protein Y with a Z-score of 14, then the S-score for the binding of that Monoclonal Antibody to protein X is equal to 29.