

9853 Pacific Heights Blvd. Suite D. San Diego, CA 92121, USA Tel: 858-263-4982

Email: info@abeomics.com

36-3143: Anti-CD62L (L-Selectin) Monoclonal Antibody(Clone: LAM1-116)

Clonality: Monoclonal Clone Name: LAM1-116 **Application:** ELISA.FACS.WB

Reactivity: Human Gene: SELL Gene ID: 6402 **Uniprot ID:** P14151

A.11; Al528707; CD62 antigen ligand; CD62L; gp90-MEL; IgA nephropathy, susceptibility to,

included; L Selectin; L-selectin; LAM1; LECAM1; LEU8; Leukocyte surface antigen Leu-8; Leukocyte-endothelial cell adhesion molecule 1; Lnhr; LSEL; Ly-22; LYAM1; Lymph node

Alternative Name: homing receptor; Lymphocyte antigen 22; Lymphocyte surface MEL-14 antigen; Pln homing

receptor; PLNHR; SELL; TQ1

Isotype: Mouse IgG1, kappa

MAb raised against supernatant from phorbol myristic acid activated human peripheral blood Immunogen Information:

leukocytes.

Description

Selectins, also designated CD62 antigens, comprise a family of carbohydrate-binding proteins involved in mediating cellular interactions with leukocytes. L-Selectin (also designated LECAM-1 or CD62L) is expressed on the majority of B and naive T cells and on most monocytes, neutrophils and eosinophils. L-Selectin interacts with specific carbohydrates expressed by activated endothelial cells. P-Selectin (also designated GMP-140 or CD62P), expressed on activated platelets and endothelial cells, and E-Selectin (also designated ELMA-1 or CD62E), expressed on endothelial cells, exhibit overlapping ligand specificities. Both recognize sialyl-Le (x) as a ligand and bind to specific carbohydrates on neutrophils and monocytes.

Product Info

Amount: 20 μg / 100 μg

200 µg/ml of Ab Purified from Bioreactor Concentrate by Protein A/G. Prepared in 10mM PBS Content:

with 0.05% BSA & 0.05% azide. Also available WITHOUT BSA & azide at 1.0mg/ml.

Antibody with azide - store at 2 to 8°C. Antibody without azide - store at -20 to -80°C. Antibody Storage condition:

is stable for 24 months. Non-hazardous.

Application Note

ELISA (For coating, order antibody without BSA); Flow Cytometry (1-2ug/million cells); Western Blot (1-2ug/ml);




Fig. 1: SDS-PAGE Analysis Purified CD62L Mouse Monoclonal Antibody (LAM1-116). Confirmation of Purity and Integrity of Antibody.