

## 36-3145: Anti-CD162 (Selectin P Ligand) Monoclonal Antibody(Clone: PSGL1/1601)

<b>Clonality :</b>	Monoclonal
<b>Clone Name :</b>	PSGL1/1601
<b>Application :</b>	IHC
<b>Reactivity :</b>	Human
<b>Gene :</b>	SELPLG
<b>Gene ID :</b>	6404
<b>Uniprot ID :</b>	Q14242
<b>Alternative Name :</b>	CD162; CLA; Cutaneous lymphocyte associated antigen; P-selectin glycoprotein ligand 1; PSGL1; Selectin P ligand; SELPLG
<b>Isotype :</b>	Mouse IgG1, kappa
<b>Immunogen Information :</b>	Recombinant human SELPLG (CD162) protein

### Description

CD162 glycoprotein functions as a high affinity counter-receptor for the cell adhesion molecules P-, E- and L- selectin expressed on myeloid cells and stimulated T lymphocytes. As such, this protein plays a critical role in leukocyte trafficking during inflammation by tethering of leukocytes to activated platelets or endothelia expressing selectins. This protein requires two post-translational modifications, tyrosine sulfation and the addition of the sialyl Lewis x tetrasaccharide (sLex) to its O-linked glycans, for its high-affinity binding activity. Aberrant expression of this gene and polymorphisms in this gene are associated with defects in the innate and adaptive immune response.

### Product Info

<b>Amount :</b>	20 µg / 100 µg
<b>Content :</b>	200 µg/ml of Ab Purified from Bioreactor Concentrate by Protein A/G. Prepared in 10mM PBS with 0.05% BSA & 0.05% azide. Also available WITHOUT BSA & azide at 1.0mg/ml.
<b>Storage condition :</b>	Antibody with azide - store at 2 to 8°C. Antibody without azide - store at -20 to -80°C. Antibody is stable for 24 months. Non-hazardous.

### Application Note

Immunohistochemistry (Formalin-fixed) (1-2ug/ml for 30 minutes at RT),(Staining of formalin-fixed tissues requires heating tissue sections in 10mM Tris with 1mM EDTA, pH 9.0, for 45 min at 95&degC followed by cooling at RT for 20 minutes);




Fig. 1: Formalin-fixed, paraffin-embedded human Spleen stained with CD162 Monoclonal Antibody (PSGL1/1601).




Fig. 2: Formalin-fixed, paraffin-embedded human Tonsil stained with CD162 Monoclonal Antibody (PSGL1/1601).