

## 36-3190: Anti-CD43 (T-Cell Marker) Monoclonal Antibody(Clone: SPN/1766R)

<b>Clonality :</b>	Monoclonal
<b>Clone Name :</b>	SPN/1766R
<b>Application :</b>	WB,FACS,IF,IHC
<b>Reactivity :</b>	Human
<b>Gene :</b>	SPN
<b>Gene ID :</b>	6693
<b>Uniprot ID :</b>	P16150
<b>Alternative Name :</b>	Galactoglycoprotein, GALGP, GPL115, Leukocyte sialoglycoprotein, Leukosialin, LSN, Sialophorin, SPN
<b>Isotype :</b>	Rabbit IgG
<b>Immunogen Information :</b>	Recombinant full-length human SPN protein

### Description

It recognizes a cell surface glycoprotein of 95/115/135kDa (depending upon the extent of glycosylation), identified as CD43. 70-90% of T-cell lymphomas and from 22-37% of B-cell lymphomas express CD43. No reactivity has been observed with reactive B-cells. So a B-lineage population that co-expresses CD43 is highly likely to be a malignant lymphoma, especially a low-grade lymphoma, rather than a reactive B-cell population. When CD43 antibody is used in combination with anti-CD20, effective immunophenotyping of the lymphomas in formalin-fixed tissues can be obtained. Co-staining of a lymphoid infiltrate with anti-CD20 and anti-CD43 argues against a reactive process and favors a diagnosis of lymphoma.

### Product Info

<b>Amount :</b>	20 µg / 100 µg
<b>Content :</b>	200 µg/ml of recombinant MAb Purified by Protein A/G. Prepared in 10mM PBS with 0.05% BSA & 0.05% azide. Also available WITHOUT BSA & azide at 1.0mg/ml.
<b>Storage condition :</b>	Antibody with azide - store at 2 to 8°C. Antibody without azide - store at -20 to -80°C. Antibody is stable for 24 months. Non-hazardous.

### Application Note

Western Blot (1-2ug/ml); Flow Cytometry (1-2ug/million cells); Immunofluorescence (1-2ug/ml); Immunohistochemistry (Formalin-fixed) (1-2ug/ml for 30 minutes at RT),(Staining of formalin-fixed tissues requires heating tissue sections in 10mM Tris with 1mM EDTA, pH 9.0, for 45 min at 95&degC followed by cooling at RT for 20 minutes);




Fig. 1: Formalin-fixed, paraffin-embedded human Tonsil stained with CD43 Rabbit Recombinant Monoclonal Antibody (SPN/1766R).




Fig. 2: SDS-PAGE Analysis Purified CD43 Rabbit Recombinant Monoclonal Antibody (SPN/1766R).




Fig. 3: Immunofluorescence Analysis of K562 cells labeling CD43 with CD43 Rabbit Recombinant Monoclonal Antibody (SPN/1766R) followed by Goat anti-Rabbit- IgG-CF488 (Green). The nuclear counterstain is NucSpot®.




Fig. 4: Western Blot Analysis of K562 cell lysate using CD43 Rabbit Recombinant Monoclonal Antibody (SPN/1766R).




Fig. 5: Flow Cytometric Analysis of PFA-fixed K562 cells using CD43 Rabbit Recombinant Monoclonal Antibody (SPN/1766R) followed by Goat anti-Rabbit- IgG-CF488 (Blue); Isotype Control (Red)