

### 36-3227: Anti-Tal1 Monoclonal Antibody(Clone: TAL1/3146)

<b>Clonality :</b>	Monoclonal
<b>Clone Name :</b>	TAL1/3146
<b>Application :</b>	ELISA
<b>Reactivity :</b>	Human
<b>Gene :</b>	TAL1
<b>Gene ID :</b>	6886
<b>Uniprot ID :</b>	P17542
<b>Alternative Name :</b>	bHLHa17; Class A basic helix-loop-helix protein 17; OTTHUMP0000009563; OTTHUMP0000009564; SCL; STEM CELL LEUKEMIA HEMATOPOIETIC TRANSCRIPTION FACTOR; Stem cell protein; T cell acute lymphocytic leukemia 1; T cell acute lymphocytic leukemia 1 protein; T cell acute lymphocytic leukemia 1 protein; T cell leukemia/lymphoma 5 protein; T-cell acute lymphocytic leukemia protein 1; T-cell leukemia/lymphoma protein 5; Tal 1 product; TAL 1 protein; TAL bHLH transcription factor 1 erythroid differentiation factor; TAL-1
<b>Isotype :</b>	Mouse IgG1, kappa
<b>Immunogen Information :</b>	Recombinant human full-length TAL1 protein

#### Description

Activation of TAL1 characterizes up to 60% of cases of human T cell acute lymphoblastic leukemia, making it the most frequent gain-of-function mutation observed in this disorder. TAL1 (also designated SCL) is a serine phosphoprotein and basic helix-loop-helix transcription factor known to regulate embryonic hematopoiesis. This transcription factor binds as a heterodimer with E2A and HEB/HTF4 to a nucleotide sequence motif termed the E-box. In addition, leukemogenesis is accelerated dramatically by transgenic co-expression of TAL1 and the catalytic subunit of casein kinase II, a serine/threonine protein kinase known to modulate the activity of other HLH transcription factors.

#### Product Info

<b>Amount :</b>	20 µg / 100 µg
<b>Content :</b>	200 µg/ml of Ab Purified from Bioreactor Concentrate by Protein A/G. Prepared in 10mM PBS with 0.05% BSA & 0.05% azide. Also available WITHOUT BSA at 1.0mg/ml.
<b>Storage condition :</b>	Antibody with azide - store at 2 to 8°C. Antibody is stable for 24 months. Non-hazardous.

#### Application Note

ELISA (For coating, order antibody without BSA)