

36-3228: Anti-TCF4 (Transcription Factor 4) Monoclonal Antibody(Clone: TCF4/1705)

Clonality :	Monoclonal
Clone Name :	TCF4/1705
Application :	ELISA,WB,IHC
Reactivity :	Human
Gene :	TCF4
Gene ID :	6925
Uniprot ID :	P15884
Alternative Name :	bHLHb19; Class B basic helix-loop-helix protein 19; Immunoglobulin transcription factor 2; ITF-2; PTHS; SEF2; SL3-3 enhancer factor 2; TCF4; Transcription factor 4
Isotype :	Mouse IgG2a, kappa
Immunogen Information :	Recombinant human TCF4 protein fragment (around aa 365-671) (exact sequence is proprietary)

Description

Recognizes a protein of 71kDa, identified as Transcription Factor 4 (TCF4). It is a basic helix-loop-helix transcription factor. The encoded protein recognizes an Ephrussi-box ('E-box') binding site ('CANNTG') - a motif first identified in immunoglobulin enhancers. This gene is broadly expressed and may play an important role in nervous system development. Defects in this gene are a cause of Pitt-Hopkins syndrome.

Product Info

Amount :	20 µg / 100 µg
Content :	200 µg/ml of Ab Purified from Bioreactor Concentrate by Protein A/G. Prepared in 10mM PBS with 0.05% BSA & 0.05% azide. Also available WITHOUT BSA & azide at 1.0mg/ml.
Storage condition :	Antibody with azide - store at 2 to 8°C. Antibody without azide - store at -20 to -80°C. Antibody is stable for 24 months. Non-hazardous.

Application Note

ELISA (For coating, order antibody without BSA); ,Western Blot (1-2ug/ml); ,Immunohistochemistry (Formalin-fixed) (1-2ug/ml for 30 minutes at RT),(Staining of formalin-fixed tissues requires heating tissue sections in 10mM Tris with 1mM EDTA, pH 9.0, for 45 min at 95 °C followed by cooling at RT for 20 minutes),




Fig. 1: Western Blot (1) HeLa and (2) HepG2 cell lysate using TCF4 Mouse Monoclonal Antibody (TCF4/1705).




Fig. 2: SDS-PAGE Analysis of Purified TCF4 Mouse Monoclonal Antibody (TCF4/1705). Confirmation of Purity and Integrity of Antibody.




Fig. 3: Analysis of Protein Array containing more than 19,000 full-length human proteins using TCF4 Mouse Monoclonal Antibody (TCF4/1705) Z- and S- Score: The Z-score represents the strength of a signal that a monoclonal antibody (Monoclonal Antibody) (in combination with a fluorescently-tagged anti-IgG secondary antibody) produces when binding to a particular protein on the HuProt™ array. Z-scores are described in units of standard deviations (SD's) above the mean value of all signals generated on that array. If targets on HuProt™ are arranged in descending order of the Z-score, the S-score is the difference (also in units of SD's) between the Z-score. S-score therefore represents the relative target specificity of a Monoclonal Antibody to its intended target. A Monoclonal Antibody is considered to specific to its intended target, if the Monoclonal Antibody has an S-score of at least 2.5. For example, if a Monoclonal Antibody binds to protein X with a Z-score of 43 and to protein Y with a Z-score of 14, then the S-score for the binding of that Monoclonal Antibody to protein X is equal to 29.