

36-3242: Anti-CD71 / Transferrin Receptor (TFRC) (Extracellular Domain) Monoclonal Antibody (Clone: TFRC/1839)

Clonality :	Monoclonal
Clone Name :	TFRC/1839
Application :	ELISA, IHC
Reactivity :	Human
Gene :	TFRC
Gene ID :	7037
Uniprot ID :	P02786
Alternative Name :	Mtvr-1, p90, TFR1, TFRC transferrin receptor (p90 CD71), TRFR
Isotype :	Mouse IgG2b, kappa
Immunogen Information :	Recombinant extracellular fragment (around aa 94-212) of human TFRC protein (exact sequence is proprietary)

Description

It recognizes a ~90-95kDa protein which is identified as cell surface transferrin receptor (CD71), a disulfide-bonded homodimeric glycoprotein of 180-190kDa. This MAb is highly specific to CD71 and shows no cross-reaction with other related proteins. Ligand for transferrin receptor is the serum iron transport protein, transferrin. This receptor is broadly distributed in carcinomas, sarcomas, leukemias, and lymphomas. CD71/Transferrin receptor has been reported to be associated with cell proliferation in both normal and neoplastic tissues and useful in predicting clinical behavior or response to therapy in a number of malignancies including breast cancer.

Product Info

Amount :	20 µg / 100 µg
Content :	200 µg/ml of Ab Purified from Bioreactor Concentrate by Protein A/G. Prepared in 10mM PBS with 0.05% BSA & 0.05% azide. Also available WITHOUT BSA & azide at 1.0mg/ml.
Storage condition :	Antibody with azide - store at 2 to 8°C. Antibody without azide - store at -20 to -80°C. Antibody is stable for 24 months. Non-hazardous.

Application Note

ELISA (Use Ab at 2-4µg/ml for coating) (Order Ab without BSA); Immunohistochemistry (Formalin-fixed) (1-2µg/ml for 30 minutes at RT)(Staining of formalin-fixed tissues requires heating tissue sections in 10mM Tris with 1mM EDTA, pH 9.0, for 45 min at 95°C followed by cooling at RT for 20 minutes);




Fig. 1: Formalin-fixed, paraffin-embedded human Placenta stained with CD71 Mouse Monoclonal Antibody (TFRC/1839).




Fig. 2: SDS-PAGE Analysis Purified CD71 Mouse Monoclonal Antibody (TFRC/1839). Confirmation of Integrity and Purity of Antibody




Fig. 3: Analysis of Protein Array containing >19,000 full-length human proteins using CD71 Mouse Monoclonal Antibody (TFRC/1839) Z- and S- Score: The Z-score represents the strength of a signal that a monoclonal antibody (Monoclonal Antibody) (in combination with a fluorescently-tagged anti-IgG secondary antibody) produces when binding to a particular protein on the HuProt™ array. Z-scores are described in units of standard deviations (SD's) above the mean value of all signals generated on that array. If targets on HuProt™ are arranged in descending order of the Z-score, the S-score is the difference (also in units of SD's) between the Z-score. S-score therefore represents the relative target specificity of a Monoclonal Antibody to its intended target. A Monoclonal Antibody is considered to specific to its intended target, if the Monoclonal Antibody has an S-score of at least 2.5. For example, if a Monoclonal Antibody binds to protein X with a Z-score of 43 and to protein Y with a Z-score of 14, then the S-score for the binding of that Monoclonal Antibody to protein X is equal to 29.