

36-3307: Anti-TPO (Thyroid Peroxidase) (Thyroid Marker) Monoclonal Antibody(Clone: TPO/3813R)

| | |
|--------------------------------|---|
| Clonality : | Monoclonal |
| Clone Name : | TPO/3813R |
| Application : | IHC |
| Reactivity : | Human |
| Gene : | TPO |
| Gene ID : | 7173 |
| Uniprot ID : | P07202 |
| Alternative Name : | MSA; TDH2A; Thyroid microsomal antigen; Thyroid peroxidase; Thyroperoxidase; TPO; TPX |
| Isotype : | Rabbit IgG |
| Immunogen Information : | Recombinant fragment of human TPO (around aa 685-804) (Exact sequence is proprietary) |

Description

Thyroperoxidase (TPO) is a 933-amino acid, type I transmembrane glycoprotein that plays a key role in thyroid gland function and autoimmunity. It is present as a dimer on the apical surface of thyroid follicular cells. TPO functions in the iodination of tyrosine residues in thyroglobulin and phenoxy-ester formation between pairs of iodinated tyrosines to generate the thyroid hormones, thyroxine and triiodothyronine. Mutations in this gene are associated with several disorders of thyroid hormonogenesis, including congenital hypothyroidism, congenital goiter, and thyroid hormone organification defect IIA. Malignant thyroid tumors exhibit an anomaly in TPO resulting in lower affinity for anti-TPO. This antibody may aid in the differentiation between benign and malignant thyroid tumors.

Product Info

| | |
|----------------------------|---|
| Amount : | 20 µg / 100 µg |
| Content : | 200 µg/ml of Ab Purified from Bioreactor Concentrate by Protein A/G. Prepared in 10mM PBS with 0.05% BSA & 0.05% azide. Also available WITHOUT BSA & azide at 1.0mg/ml. |
| Storage condition : | Antibody with azide - store at 2 to 8°C. Antibody without azide - store at -20 to -80°C. Antibody is stable for 24 months. Non-hazardous. |

Application Note

Immunohistochemistry (Formalin-fixed) (1-2ug/ml for 30 minutes at RT),(Staining of formalin-fixed tissues requires heating tissue sections in 10mM Tris with 1mM EDTA, pH 9.0, for 45 min at 95°C followed by cooling at RT for 20 minutes);




Fig. 1: SDS-PAGE Analysis Purified TPO Recombinant Rabbit Monoclonal Antibody (TPO/3813R). Confirmation of Integrity and Purity of the Antibody.